

Vacua of Massive Hyper-Kähler Sigma Models with Non-Abelian Quotient

Masato Arai^{1,*}, Muneto Nitta^{2,*} and Norisuke Sakai³

¹*Institute of Physics, AS CR, 182 21, Praha 8, Czech Republic*

²*Department of Physics, Purdue University, West Lafayette, IN 47907-1396, USA*

³*Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan*

The Higgs branch of $\mathcal{N} = 2$ supersymmetric gauge theories with non-Abelian gauge groups are described by hyper-Kähler (HK) nonlinear sigma models with potential terms. Using the non-Abelian HK quotient with respect to $U(M)$ and $SU(M)$ gauge groups, we derive the massive HK sigma models that are not toric in the $\mathcal{N} = 1$ superfield formalism and the harmonic superspace formalism. The $U(M)$ quotient gives $N!/[M!(N-M)!]$ discrete vacua that may allow various types of domain walls, whereas the $SU(M)$ quotient gives no discrete vacua.

§1. Introduction

Possibilities of large extra dimensions^{1),2)} has renewed interest in field theories in higher-dimensional systems. In this brane-world scenario, our four-dimensional world is realized on an extended topological defect such as a wall (brane). Supersymmetry (SUSY) has been an extremely useful principle for building unified models beyond the standard model.³⁾ In SUSY theories, topological defects can often be obtained as BPS states,⁴⁾ which preserve part of the SUSY.⁵⁾ The BPS states play an important role in exploring nonperturbative effects in SUSY gauge theories.⁶⁾ The Higgs branch of the SUSY gauge theories can often be described by SUSY nonlinear sigma models.^{7),8)} In turn, these SUSY nonlinear sigma models are useful for obtaining topological defects, such as domain walls and flux tubes.^{9),10)} Supersymmetric theories in systems of greater than four dimensions require at least eight supercharges. The scalar and spinor matter fields can be described by means of hypermultiplets in theories with eight SUSY. Recently, we formulated $\frac{1}{2}$ single-BPS

* Present address: Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan.

domain walls in an eight SUSY model in four dimensions.¹¹⁾ Moreover, we have also succeeded in constructing the $\frac{1}{2}$ BPS wall¹²⁾ and BPS multi-walls¹³⁾ consistently in five-dimensional supergravity. Before discussing the SUSY five-dimensional theories, it is useful to consider models with eight SUSY in four dimensions without gravity.

Theories with eight SUSY are so restrictive that the nontrivial interactions require the nonlinearity of the kinetic term (the nonlinear sigma model) if there are only hypermultiplets. Target manifolds of the SUSY nonlinear sigma models with eight SUSY must be hyper-Kähler (HK) manifolds.^{14),15)} To obtain a wall solution, we need a nontrivial potential. The potential term for any HK sigma model is severely restricted by eight SUSY.¹⁶⁾ It can be written as the square of the triholomorphic Killing vector in the on-shell formulation, and it can be understood in terms of the Scherk-Schwarz reduction¹⁷⁾ from six dimensions.¹⁸⁾ These models are called “massive HK nonlinear sigma models”. Contrastingly, in two dimensions, $\mathcal{N} = 2$ SUSY (four SUSY) sigma models can have a similar potential term,¹⁶⁾ in addition to the potential that can be derived from the superpotential and can be constructed off-shell in the superfield formalism.¹⁹⁾ Solitons in such models are discussed in Ref. 20). The off-shell formulation of $\mathcal{N} = 2$ massive HK sigma models in four dimensions was given recently.¹¹⁾ An exact BPS solution of the wall junction was also constructed recently in an $\mathcal{N} = 2$ theory with hypermultiplets and a vector multiplet.²¹⁾

A large class of the HK manifold is given by toric HK manifolds that are defined as HK manifolds of real dimension $4n$ admitting n mutually commuting Abelian triholomorphic isometries.^{22)–25)} In Ref. 26), the massive HK sigma model on any toric HK manifold was derived in the component formalism in four-dimensional spacetime. The solution of the N parallel domain walls was found for $T^*\mathbb{CP}^{N-1}$, which is a toric HK manifold, and its moduli space was constructed.²⁷⁾ For this massive $T^*\mathbb{CP}^{N-1}$ model, the dynamics of the walls were studied²⁸⁾ and the number of zero modes were calculated using the index theorem.²⁹⁾ The off-shell formulation of the massive $T^*\mathbb{CP}^{N-1}$ model in four-dimensional spacetime was obtained for both $\mathcal{N} = 1$ superfields and $\mathcal{N} = 2$ superfields in harmonic superspace formalism (HSF).¹¹⁾ Other interesting solitons, like solutions representing strings ending on walls, wall intersections and string intersections were also considered in the toric HK nonlinear

sigma models.^{26),30)–33)}

The potential term of the massive $T^*\mathbf{CP}^{N-1}$ model comes from the mass terms of the hypermultiplets when the nonlinear sigma model is constructed as the quotient with respect to the $U(1)$ gauge group.^{11),28)} We call this formulation of massive HK sigma models “the massive HK quotient method,” since the massless case is identified to a HK quotient found in Refs. 22) and 35). One of the advantages of our massive HK quotient is that the off-shell formulation of the SUSY nonlinear sigma models is possible.¹¹⁾ The off-shell formulation is powerful for extending models to those with other isometries and/or gauge symmetries and to those coupled to gravity, because in it (part of) SUSY is manifest. Any *toric* HK manifold can be constructed using an *Abelian* HK quotient.^{24),25)} Therefore, an off-shell formulation of general massive toric HK sigma models²⁶⁾ can be obtained using the massive HK quotient with Abelian gauge theories. By contrast, a HK nonlinear sigma model other than the toric HK target manifolds has been obtained by Lindström and Roček²²⁾ as a quotient using a non-Abelian gauge group for the *massless* case only (without potential terms).

The purpose of this paper is to investigate *massive* HK sigma models in four-dimensional spacetime using the quotient of the SUSY QCD with respect to a *non-Abelian* gauge group. The models are no longer toric HK manifolds, and turn out to be the cotangent bundle over the complex Grassmann manifold $T^*G_{N,M}$ and its generalization. The former model is the massive generalization of the massless nonlinear sigma model presented in Ref. 22). We obtain potential terms for this massive HK nonlinear sigma model and investigate the vacuum structure of the massive $T^*G_{N,M}$ model in detail. We find that this model has $N!/ [M!(N-M)!]$ discrete vacua which are characterized by mutually orthogonal M -dimensional complex hyperplanes in $\mathbf{C}^N$. We therefore can expect rich structure for wall solutions in this model. By considering the $SU(M)$ gauge group, we obtain a generalization of the massive $T^*G_{N,M}$ model, the massive HK nonlinear sigma model with the quaternionic line bundle over the $T^*G_{N,M}$ as the target space. The vacua of this model are trivial, and hence we cannot expect interesting solitons.

There exists a similar model with the same number of vacua, namely the two-dimensional $\mathcal{N} = 2$ SUSY (four SUSY) Grassmann $G_{N,M}$ sigma model with twisted

mass.³⁴⁾ We believe that the target space of this model is obtained by truncating our model on $T^*G_{N,M}$ to its base manifold, so that they have the same number of vacua. If we find solitons in our model, it may also be the case that they play interesting roles in this kind of two dimensional model.

The rest of this paper is organized as follows. In §2, we present the massive HK quotient with $\mathcal{N} = 1$ superfields. The vacuum structure of the massive $T^*G_{N,M}$ model is studied in §3. The massive HK quotient is described in the Wess-Zumino gauge in §4. Section 5 is devoted to a generalization of the massive HK quotient with respect to the SU gauge group. Our model is derived in the harmonic superspace formalism^{36),37)} in §6. A summary and discussion are given in §7.

§2. Massive HK quotient with respect to the $U(M)$ gauge group

We consider $\mathcal{N} = 2$ SUSY QCD with N flavors and a $U(M)$ gauge group. In terms of $\mathcal{N} = 1$ superfields, the NM $\mathcal{N} = 2$ hypermultiplets ** can be decomposed into $(N \times M)$ - and $(M \times N)$ -matrix chiral superfields $\Phi(x, \theta, \bar{\theta})$ and $\Psi(x, \theta, \bar{\theta})$, and the $\mathcal{N} = 2$ vector multiplets for the $U(M)$ gauge symmetry can be decomposed into $M \times M$ matrices of the $\mathcal{N} = 1$ vector superfields $V = V^A(x, \theta, \bar{\theta})T_A$ and chiral superfields $\Sigma = \Sigma^A(x, \theta, \bar{\theta})T_A$, with the $M \times M$ matrices T_A ($A = 1, \dots, M^2$) of the fundamental representation of the generators of the $U(M)$ gauge group. The $U(M)$ gauge transformation is given by

$$e^V \rightarrow e^{V'} = e^{-i\Lambda} e^V e^{i\Lambda^\dagger}, \quad \Sigma \rightarrow \Sigma' = e^{-i\Lambda} \Sigma e^{i\Lambda}, \quad (2.1)$$

$$\Phi \rightarrow \Phi' = \Phi e^{i\Lambda}, \quad \Psi \rightarrow \Psi' = e^{-i\Lambda} \Psi, \quad (2.2)$$

where $\Lambda = \Lambda^A(x, \theta, \bar{\theta})T_A$ are the chiral superfields of the gauge parameters. Note that this gauge symmetry is actually complexified into $U(M)^{\mathbf{C}} = GL(M, \mathbf{C})$, because the scalar components of Λ are complex fields.

Because we focus on the Higgs branch, we can take the strong coupling limit, $g \rightarrow \infty$, and throw away the kinetic term for gauge multiplets. Then, the Lagrangian

** In the following, we consider only the case $N > M$. This requirement is necessary in order for the nonlinear sigma models to be well-defined, since dimensions of target space is given by $4(N - M)$.

is given by

$$\begin{aligned} \mathcal{L} = & \int d^4\theta \left[\text{tr}(\Phi^\dagger \Phi e^V) + \text{tr}(\Psi \Psi^\dagger e^{-V}) - c \text{tr} V \right] \\ & + \left[\int d^2\theta \left(\text{tr} \{ \Sigma (\Psi \Phi - b \mathbf{1}_M) \} + \sum_{a=1}^{N-1} m_a \text{tr}(\Psi H_a \Phi) \right) + \text{c.c.} \right], \end{aligned} \quad (2.3)$$

where we have absorbed a common hypermultiplet mass into the field Σ , the quantities m_a ($a = 1, \dots, N-1$) are complex mass parameters, and H_a are diagonal traceless matrices, interpreted as the Cartan generators of $SU(N)$ below. The electric and magnetic Fayet-Iliopoulos (FI) parameters are denoted $c \in \mathbf{R}_{\geq 0}$ and $b \in \mathbf{C}$, respectively. In the limit $m_a \rightarrow 0$ for all a , the model has the global (flavor) $SU(N)$ symmetry expressed by

$$\Phi \rightarrow \Phi' = g\Phi, \quad \Psi \rightarrow \Psi' = \Psi g^{-1}, \quad g \in SU(N), \quad (2.4)$$

with V and Σ unchanged.^{***} For nonzero m_a , this $SU(N)$ symmetry is explicitly broken down to $U(1)^{N-1}$, generated by H_a .

Setting $\Psi = 0$ in the Lagrangian (2.3), it reduces to the Kähler quotient construction of the (complex) Grassmann manifold $G_{N,M} = SU(N)/[SU(M) \times SU(N-M) \times U(1)]$,³⁸⁾ and this suggests that the full Lagrangian is related to $G_{N,M}$. Actually, for the massless case, this Lagrangian reduces to the HK quotient construction of $T^*G_{N,M}$,²²⁾ generalizing the $U(1)$ HK quotient construction^{39)–41)} for $T^*\mathbf{C}P^{N-1}$ ($M = 1$) with the Calabi metric.⁴²⁾

Since we have introduced the mass parameter through the H_a generator, we will eventually obtain a potential term that is the square of the tri-holomorphic Killing vector corresponding to a linear combination of the matrices H_a , after eliminating the vector multiplet.

Next, we eliminate the auxiliary superfields V and Σ in the superfield formalism. From Eq. (2.3), we find that their equations of motion read [†]

$$\frac{\partial \mathcal{L}}{\partial V} = \Phi^\dagger \Phi e^V - e^{-V} \Psi \Psi^\dagger - c \mathbf{1}_M = 0, \quad (2.5)$$

$$\frac{\partial \mathcal{L}}{\partial \Sigma} = \Psi \Phi - b \mathbf{1}_M = 0. \quad (2.6)$$

^{***} When $b = 0$, there exists an additional $U(1)$ given by $\Phi \rightarrow \Phi' = e^{i\alpha} \Phi, \Psi \rightarrow \Psi' = e^{i\alpha} \Psi, \Sigma \rightarrow \Sigma' = e^{-2i\alpha} \Sigma$, but this is inconsistent with $\mathcal{N} = 2$ SUSY.

[†] Here we take $\delta X \equiv e^{-V} \delta e^V$ as an infinitesimal parameter of variation. Then, the equations $\delta \text{tr}(\Phi^\dagger \Phi e^V) = \text{tr}(\Phi^\dagger \Phi e^V \delta X)$ and $\delta \text{tr}(\Psi \Psi^\dagger e^{-V}) = -\text{tr}(e^{-V} \Psi \Psi^\dagger \delta X)$ hold.

Multiplying Eq. (2.5) by $\Phi^\dagger \Phi e^V$ from the left, we find

$$\left(\Phi^\dagger \Phi e^V - \frac{c}{2} \mathbf{1}_M\right)^2 - \frac{c^2}{4} \mathbf{1}_M - \Phi^\dagger \Phi \Psi \Psi^\dagger = 0. \quad (2.7)$$

From the above equation, V can be easily solved, and we have

$$e^V = \frac{c}{2} (\Phi^\dagger \Phi)^{-1} \left(\mathbf{1}_M \pm \sqrt{\mathbf{1}_M + \frac{4}{c^2} \Phi^\dagger \Phi \Psi \Psi^\dagger} \right). \quad (2.8)$$

Substituting this back into (2.3), we obtain the Kähler potential for the Lindström-Roček metric,²²⁾

$$K = c \operatorname{tr} \sqrt{\mathbf{1}_M + \frac{4}{c^2} \Phi^\dagger \Phi \Psi \Psi^\dagger} - c \operatorname{tr} \log \left(\mathbf{1}_M + \sqrt{\mathbf{1}_M + \frac{4}{c^2} \Phi^\dagger \Phi \Psi \Psi^\dagger} \right) + c \operatorname{tr} \log \Phi^\dagger \Phi, \quad (2.9)$$

where we have used the formula $\operatorname{tr} \log AB = \operatorname{tr} \log A + \operatorname{tr} \log B$, which holds for any square matrices A and B , and we have omitted constant terms because they vanish under the superspace integration. Here we have chosen the plus sign in Eq. (2.8) to realize the positivity of the metric. Using the $N \times N$ meson matrix of the gauge invariant,

$$\mathcal{M} = \Phi \Psi, \quad (2.10)$$

the Kähler potential can be rewritten as that in Ref. 8),

$$K = c \frac{M}{N} \operatorname{tr}_{N \times N} \sqrt{\mathbf{1}_N + \frac{N}{M} \frac{4}{c^2} \mathcal{M} \mathcal{M}^\dagger} - c \frac{M}{N} \operatorname{tr}_{N \times N} \log \left(\mathbf{1}_N + \sqrt{\mathbf{1}_N + \frac{N}{M} \frac{4}{c^2} \mathcal{M} \mathcal{M}^\dagger} \right) + c \operatorname{tr} \log \Phi^\dagger \Phi, \quad (2.11)$$

where we have made explicit that the traces are taken in $N \times N$ matrices, whereas the previous ones $M \times M$.

The Kähler potential (2.9) or (2.11) is strictly invariant under the full global (flavor) $SU(N)$ symmetry (2.4), because the mass term in the Lagrangian (2.3) does not affect the D-term. In Eqs. (2.9) and (2.11), the first two terms are strictly invariant under the gauge transformation (2.2) for the matter fields, but the third term induces the Kähler transformation

$$K \rightarrow K' = K + ic \operatorname{tr} \Lambda - ic \operatorname{tr} \Lambda^\dagger. \quad (2.12)$$

Fixing this complexified $U(M)$ gauge symmetry and solving the constraint (2.6), we obtain the Lagrangian of the nonlinear sigma model in terms of independent superfields. For this purpose, we should consider the two cases $b = 0$ and $b \neq 0$ separately. Note that the physics does not depend on this choice, as the two cases are transformed into each other under the $SU(2)_R$ transformation. Although this transformation does not preserve the holomorphy, these two cases merely involve different complex structures.

i) $b = 0$. In this case, the gauge can be fixed as

$$\Phi = \begin{pmatrix} \mathbf{1}_M \\ \varphi \end{pmatrix}, \quad \Psi = (-\psi\varphi, \psi), \quad (2.13)$$

with φ and ψ being $[(N-M) \times M]$ - and $[M \times (N-M)]$ -matrix chiral superfields, respectively. Then, the superpotential becomes

$$W = \sum_a m_a \text{tr} \left[(-\psi\varphi, \psi) H_a \begin{pmatrix} \mathbf{1}_M \\ \varphi \end{pmatrix} \right] = \sum_a m_a \text{tr} \left[H_a \begin{pmatrix} -\psi\varphi & \psi \\ -\varphi\psi\varphi & \varphi\psi \end{pmatrix} \right]. \quad (2.14)$$

ii) $b \neq 0$. In this case, we can take the gauge as²²⁾

$$\Phi = \begin{pmatrix} \mathbf{1}_M \\ \varphi \end{pmatrix} Q, \quad \Psi = Q(\mathbf{1}_M, \psi), \quad Q = \sqrt{b}(\mathbf{1}_M + \psi\varphi)^{-\frac{1}{2}}, \quad (2.15)$$

with φ and ψ again being $[(N-M) \times M]$ - and $[M \times (N-M)]$ -matrix chiral superfields, respectively. In this case, the superpotential is given by

$$W = b \sum_a m_a \text{tr} \left[H_a \begin{pmatrix} \mathbf{1}_M \\ \varphi \end{pmatrix} (\mathbf{1}_M + \psi\varphi)^{-1} (\mathbf{1}_M, \psi) \right]. \quad (2.16)$$

We can find the bundle structure of the manifold as follows. First, let us consider the $b = 0$ case. Then, setting $\psi = 0$, the Kähler potential becomes

$$K|_{\psi=0} = c \text{tr} \log(1 + \varphi^\dagger \varphi), \quad (2.17)$$

which is that of the Grassmann manifold. Therefore, φ parameterizes the base Grassmann manifold, whereas ψ parameterizes the cotangent space as the fiber,²²⁾ with the total space being the cotangent bundle over the Grassmann manifold $T^*G_{N,M}$.

Next, consider the case $b \neq 0$. In the case of $T^*\mathbf{CP}^{N-1}$ with $M = 1$, the base manifold is embedded by $\varphi = \psi^\dagger$.⁴⁰⁾††

The tri-holomorphic isometry group of this manifold is $G = SU(N)$, whereas the isotropy group is $H = SU(M) \times SU(N-M) \times U(1)$, since the manifold is $T^*(G/H)$. In each case $b = 0$ and $b \neq 0$, the action of $g \in SU(N)$ is found to be as follows:

$$\Phi \rightarrow \Phi' = g\Phi h^{-1}(\varphi, \psi), \quad \Psi \rightarrow \Psi' = h(\varphi, \psi)\Psi g^{-1}. \quad (2.18)$$

Here, $h(\varphi, \psi) = e^{-i\Lambda(\varphi, \psi)}$ is an induced gauge transformation called a 'compensator'. This is needed to put $g\Phi$ and Ψg^{-1} back into the gauge slice, because they are not in general in the gauge fixing condition (2.13) or (2.15). This makes the (base) manifold a coset space G/H . Under the global $SU(N)$ transformation, the Kähler potential undergoes the Kähler transformation

$$\begin{aligned} K \rightarrow K' &= K + \text{ctr log}(h^{-1}(\varphi, \psi))^\dagger + \text{ctr log } h^{-1}(\varphi, \psi) \\ &= K + ic \text{tr } \Lambda(\varphi, \psi) - ic \text{tr } \Lambda^\dagger(\varphi^\dagger, \psi^\dagger), \end{aligned} \quad (2.19)$$

whereas the Kähler metric is invariant. The tri-holomorphic Killing vectors for the isometry $SU(N)$ can be calculated from infinitesimal transformations of (2.18).

Here we give several more comments.

1. There exists a manifest duality between the two theories with the $U(M)$ and $U(N-M)$ gauge symmetries and the same flavor $SU(N)$ symmetry. This results directly from the duality in the base Grassmann manifold $G_{N,M} \simeq G_{N,N-M}$.
2. For $M = 1$ ($M = N-1$), namely for the $U(1)$ [$U(N-1)$] gauge symmetry, this model reduces to $T^*\mathbf{CP}^{N-1} \simeq T^*G_{N,1} (\simeq T^*G_{N,N-1})$,⁴²⁾ which is discussed in detail in Ref. 11). Moreover, if $N = 2$, the manifold $T^*\mathbf{CP}^1$ is the Eguchi-Hanson space.⁴³⁾

There are nontrivial models that do not reduce to $T^*\mathbf{CP}^{N-1}$ target manifold. The nontrivial model with the lowest dimensions of the target manifold is the case with $N = 4, M = 2$. In this case, the manifold is $T^*G_{4,2} = T^*[SU(4)/SU(2) \times SU(2) \times U(1)] = T^*[SO(6)/SO(4) \times U(1)] \equiv T^*Q^4$, in which the base manifold Q^4 is called the Klein quadric space.^{†††}

†† The embedding $\varphi = \psi^\dagger$ should hold for a matrix with any value of M , although we have not yet proved this.

††† For diffeomorphisms of base manifolds, see Ref. 44).

§3. Vacuum structure

In the massless case, $m_a = 0$, the moduli space of the vacua is $T^*G_{N,M}$ itself (the Higgs branch of $\mathcal{N} = 2$ SUSY QCD^{(7), (8)}), since the low energy effective theory has no potential term. However, once the mass term is switched on, most vacua are lifted up, leaving some discrete points as vacua. First we consider the simpler case of $T^*\mathbf{CP}^{N-1}$ and then that of $T^*G_{N,M}$.

3.1. Vacua in the massive $T^*\mathbf{CP}^{N-1}$ model

In this subsection, we investigate $T^*\mathbf{CP}^{N-1} = T^*G_{N,1}$ with $M = 1$. Without loss of generality we consider the case $b = 0$ with $c \neq 0$. In this case, the dynamical matrix fields are column and row vectors like $\varphi^T = (\varphi^1, \dots, \varphi^{N-1})$ and $\psi = (\psi^1, \dots, \psi^{N-1})$.

The superpotential given in (2.14) becomes

$$W = \sum_a m_a \text{tr} \left[H_a \begin{pmatrix} -\psi \cdot \varphi & \psi \\ -\varphi(\psi \cdot \varphi) & \varphi \otimes \psi \end{pmatrix} \right]. \quad (3.1)$$

We choose H_a ($a = 1, \dots, N-1$) as

$$H_a = \frac{1}{\sqrt{a(a+1)}} \text{diag.} (1, \dots, 1, -a, 0, \dots, 0), \quad (3.2)$$

where $-a$ is the $(a+1)$ -th component, with the normalization given by the trace $\text{tr}(H_a H_b) = \delta_{ab}$. Then the superpotential can be calculated as

$$W = - \sum_a M_a \psi^a \varphi^a, \quad M_a \equiv \sqrt{\frac{a}{a+1}} m_a + \sum_{b=1}^a \frac{m_b}{\sqrt{b(b+1)}}. \quad (3.3)$$

Therefore the derivatives of W with respect to the fields are

$$\partial_{\varphi^a} W = -M_a \psi^a, \quad \partial_{\psi^a} W = -M_a \varphi^a \quad (\text{no sum}). \quad (3.4)$$

These vanish only at the origin, $\varphi = \psi^T = 0$, provided that $M_a \neq 0$. This is the only vacuum in the regular region of these coordinates, because the metric is regular there.

However, this model contains more vacua, because the entire manifold is covered by several coordinate patches, and a vacuum exists at the origin of each coordinate patch. To see this, we temporarily concentrate on the base $\mathbf{CP}^{N-1}$. We consider the

fields before the gauge fixing, $\Phi \equiv \phi^A = (\phi^1, \dots, \phi^N)^T$ ($A = 1, \dots, N$), called the homogeneous coordinates, in which we need an identification by the gauge transformation $\phi^A \sim e^{i\Lambda} \phi^A$ to realize the space $\mathbf{CP}^{N-1}$. In the region satisfying $\phi^1 \neq 0$, we can choose a patch $\varphi^i = \phi^{i+1}/\phi^1$ ($i = 1, \dots, N-1$), which was used in Eq. (2.13). Here, let us write these coordinates as $\varphi_{(1)}^i = \phi^{i+1}/\phi^1$. In the same way, in the region satisfying $\phi^A \neq 0$, we can choose the A -th patch defined by

$$\varphi_{(A)}^i = \begin{cases} \phi^i/\phi^A, & (1 \leq i \leq A-1) \\ \phi^{i+1}/\phi^A. & (A \leq i \leq N-1) \end{cases}. \quad (3.5)$$

We thus have N sets of patches $\{\varphi_{(A)}^i\}$, which is sufficient to cover the whole base manifold. Corresponding to each patch for the base space, we manifestly have an associated patch for the fiber tangent space $\{\psi_{(A)}^i\}$, from Eq. (2.13). These sets of coordinates, $\{\varphi_{(A)}^i, \psi_{(A)}^i\}$, are sufficient to cover the whole $T^*\mathbf{CP}^{N-1}$. For each patch, the origin $\varphi_{(A)}^i = \psi_{(A)}^i = 0$ is a vacuum. Therefore, the number of discrete vacua for the massive $T^*\mathbf{CP}^{N-1}$ model is N . This result was first obtained in Ref. 27).

To investigate solitons like BPS walls, their junction and lumps, it may be better to consider the problem in one coordinate patch. The other vacua appear in one patch as the coordinate singularities of the metric in infinities of the coordinates rather than the stationary points of the superpotential.⁴⁵⁾ To see this, we again consider only the base $\mathbf{CP}^{N-1}$. We study how the A -th vacuum ($A \neq 1$) in the origin of the A -th coordinate patch is mapped into the first patch. The A -th vacuum is represented by $\varphi_{(A)}^i = 0$ or $\phi^B/\phi^A = 0$ for all B ($\neq A$). In the first coordinate patch, this point is mapped to an infinite point represented by

$$\varphi_{(1)}^{A-1} \rightarrow \infty, \quad \varphi_{(1)}^i/\varphi_{(1)}^{A-1} \rightarrow 0 \quad (i \neq A-1), \quad (3.6)$$

which appears to be a runaway vacuum in this patch. Hence, the origin and $N-1$ infinities are vacua in each coordinate patch.⁴⁵⁾ To summarize, if we include runaway vacua, one patch is sufficient to describe soliton solutions.[‡] However, note that the terminology “runaway” is merely a coordinate-dependent concept, because a

[‡] The domain wall solution connects the origin and the infinity in this parameterization for the $T^*\mathbf{CP}^1$ case.¹¹⁾

runaway vacuum in one coordinate patch is a true vacuum in another coordinate patch.


We can also study the vacua without referring to the local coordinate patches. We concentrate on the base $\mathbf{CP}^{N-1}$ once again. A point in $\mathbf{CP}^{N-1}$ corresponds to a complex line through the origin in $\mathbf{C}^N$ with homogeneous coordinates ϕ^A , because the gauge transformation corresponds to the equivalence relation $\phi^A \sim e^{iA}\phi^A$. The first vacuum is expressed in the region satisfying $\phi^1 \neq 0$ by the relations $\varphi_{(1)}^i = \phi^{i+1}/\phi^1 = 0$ ($i = 1, \dots, N-1$), namely $\phi^{i+1} = 0$. Therefore, the first vacuum corresponds to the ϕ^1 -axis. Similarly, the A -th vacuum corresponds to the ϕ^A -axis. In this way, each vacuum is simply expressed by one orthogonal axis in $\mathbf{C}^N$. Note that each axis is invariant under the $U(1)^{N-1}$ transformation of H_a , and hence it is a fixed point of this transformation.

If we take N orthogonal normalized basis vectors e_A [with $(e_A)^* \cdot e_B = \delta_{AB}$], whose components are given by

$$(e_A)^B = \delta_A^B, \quad (3.7)$$

then for any complex line in $\mathbf{C}^N$, there exists a unit vector $e' = \sum_{A=1}^N a^A e_A = U e_1$ that spans this line, where each a^A is a complex number with $\sum_A |a^A|^2 = 1$ and U is a unitary matrix, $U \in U(N)$. Each of the N -vacua found above corresponds to a different one of the vectors e_A ($A = 1, \dots, N$) (with a zero value of the cotangent space $\psi = 0$).

Example: the Eguchi-Hanson space.⁴³⁾ The simplest model is the Eguchi-Hanson space, $T^*\mathbf{CP}^1$ ($N = 2, M = 1$). This model has two discrete vacua and admits a typical domain wall solution.^{11), 30)} The vacua are located at the north and south poles of the base $\mathbf{CP}^1 \simeq S^2$ (see Fig. 1). Corresponding to the two gauge fixing conditions $\Phi = \begin{pmatrix} 1 \\ z \end{pmatrix}$ and $\Phi = \begin{pmatrix} w \\ 1 \end{pmatrix}$, we have the two coordinate patches $z \equiv \varphi_{(1)}^1 = \phi^2/\phi^1$ and $w \equiv \varphi_{(2)}^1 = \phi^1/\phi^2$, where we have $z = 1/w$. Two vacua are given by $z = 0$ and $w = 0$. The second (first) vacuum, $w = 0$ ($z = 0$), is mapped to $z = \infty$ ($w = \infty$) in the first (second) patch, which looks like a runaway vacuum. In homogeneous coordinates, these correspond to $\langle \Phi \rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \equiv e_1$ and $\langle \Phi \rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \equiv e_2$, respec-

Fig. 1. The base manifold of $T^*\mathbf{CP}^1$ and vacua.

Corresponding to two gauge fixing conditions, we have two coordinates, z and w , covering S^2 , except for the south (S) and north (N) poles, respectively. The origins of z and w (N and S, respectively) are both vacua. The domain wall solution, approaching these two vacua at spatial infinities, is mapped to a trajectory connecting N and S in S^2 .

tively, with $\langle \Psi \rangle = (0, 0)$. Also, in a coordinate independent manner, these two vacua correspond to the ϕ^1 and ϕ^2 axes, spanned by e_1 and e_2 , respectively.

Before closing this subsection, we consider the case $b \neq 0$. In this case, The superpotential (2.16) can be calculated, and we have

$$W = \frac{b}{1 + \psi \cdot \varphi} \left(L + \sum_{a=1}^{N-1} N_a \psi^a \varphi^a \right),$$

$$L \equiv \sum_{a=1}^{N-1} \frac{m_a}{\sqrt{a(a+1)}}, \quad N_a \equiv -\sqrt{\frac{a}{a+1}} m_a + \sum_{b=a+1}^{N-1} \frac{m_b}{\sqrt{b(b+1)}} = L - M_a, \quad (3.8)$$

with M_a defined in (3.3). The derivatives of W are given by

$$\partial_{\varphi^a} W = -\frac{b\psi^a}{(1 + \psi \cdot \varphi)^2} \left[L - N_a + \sum_{b=1}^{N-1} (N_b - N_a) \psi^b \varphi^b \right]$$

$$\begin{aligned}
 &= -\frac{b\psi^a}{(1 + \psi \cdot \varphi)^2} \left[M_a - \sum_{b=1}^{N-1} (M_b - M_a) \psi^b \varphi^b \right], \\
 \partial_{\psi^a} W &= (\psi^a \leftrightarrow \varphi^a), \tag{3.9}
 \end{aligned}$$

where the arrow in the second equation indicates the exchange of quantities in the first equation. The origin, $\varphi^a = \psi^a = 0$, in each patch is a vacuum. There is no vacuum other than these N vacua. The number of vacua should coincide with that in the case with $b = 0$ and $c \neq 0$, because they are connected by the R-symmetry, and the physics does not depend on the difference.

3.2. Vacua in the massive $T^*G_{N,M}$ model

To determine the vacua of the $T^*G_{N,M}$ model, we consider the case $b = 0$ and $c \neq 0$, again without loss of generality. We write the matrix φ as $(\varphi_{i\alpha})$ and ψ as $(\psi_{\alpha i})$, with the indices $i = 1, \dots, N - M$ and $\alpha = 1, \dots, M$. The superpotential given in Eq. (2.14) can be calculated as

$$\begin{aligned}
 W &= -\sum_{\alpha=1}^M \sum_{i=1}^{N-M} M_{\alpha i} \varphi_{i\alpha} \psi_{\alpha i}, \\
 M_{\alpha i} &\equiv \sqrt{\frac{i+M-1}{i+M}} m_{i+M-1} - \sqrt{\frac{\alpha-1}{\alpha}} m_{\alpha-1} + \sum_{a=\alpha}^{i+M-1} \frac{m_a}{\sqrt{a(a+1)}}, \tag{3.10}
 \end{aligned}$$

where we have set $m_0 \equiv 0$. For the case $M = 1$ ($\alpha = 1$), this reduces to Eq. (3.3) for $T^*\mathbf{CP}^{N-1}$. The derivatives of superpotential given in (3.10) with respect to the fields are

$$\partial_{\varphi_{i\alpha}} W = -M_{\alpha i} \psi_{\alpha i}, \quad \partial_{\psi_{\alpha i}} W = -M_{\alpha i} \varphi_{i\alpha} \quad (\text{no sum}). \tag{3.11}$$

Therefore, the origin of these coordinates, $\varphi = \psi^T = 0$, is a vacuum, provided that $M_{ai} \neq 0$. This is the unique vacuum in the finite region of these coordinates where the metric is regular. The number of vacua in this model is equal to the number of coordinate patches, as in the $T^*\mathbf{CP}^{N-1}$ case. In the first coordinate patch, we have chosen the first M row vectors in Φ as the unit matrix, as in Eqs. (2.13) or (2.15). The other coordinate patches are obtained by making other choices of the gauge fixing conditions, with which the other sets of M row vectors in Φ form the unit matrix. The number of such coordinate systems is ${}_N C_M = N!/[M!(N-M)!]$. They are independent and sufficient to cover the entire manifold. Therefore, this model

has $N!/([M!(N-M)!])$ vacua. This number is invariant under the duality between $U(M)$ and $U(N-M)$ gauge groups. It also reduces correctly to N for $T^*\mathbf{CP}^{N-1}$ when $M=1$ or $M=N-1$.

As in the $T^*\mathbf{CP}^{N-1}$ case, we can understand the vacua of $T^*G_{N,M}$ without local coordinates. A point in the base $G_{N,M}$ corresponds to an M -dimensional complex hyperplane (M -plane) through the origin in $\mathbf{C}^N$. The vacua found above correspond to mutually orthogonal M -planes spanned by M arbitrary sets of axes chosen from the N axes. Therefore, the total number of vacua is ${}_NC_M = N!/([M!(N-M)!])$. Because the M -planes of vacua are invariant under $U(1)^{N-1}$ generated by H_a , the vacua are fixed points.

Choosing the basis (3.7) in $\mathbf{C}^N$, a point in $G_{N,M}$ expressed as an M -plane in $\mathbf{C}^N$ can be spanned by a set of M unit vectors,

$$(e_i)' = Ue_i, \quad (3.12)$$

where $i = 1, \dots, N-M$ and U is a unitary matrix, $U \in U(N)$. The vacua of mutually orthogonal M -planes are spanned by M arbitrary sets of basis vectors among the orthogonal N -basis.

The duality becomes manifest in this framework. We can represent a point in $G_{N,M}$ by an $(N-M)$ -plane that is the complement of an M -plane.

Example: the cotangent bundle over the Klein quadric. We now consider the example of the Klein quadric $T^*G_{4,2} = T^*Q^4$ ($N=4$ and $M=2$). In this case, there exist six coordinate systems $\varphi_{i\alpha}^{(A)}$ ($A=1, \dots, 6$) for the base manifold corresponding to six choices of gauge fixing, given by

$$\Phi = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ \varphi_{11}^{(1)} & \varphi_{12}^{(1)} \\ \varphi_{21}^{(1)} & \varphi_{22}^{(1)} \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ \varphi_{11}^{(2)} & \varphi_{12}^{(2)} \\ 0 & 1 \\ \varphi_{21}^{(2)} & \varphi_{22}^{(2)} \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ \varphi_{11}^{(3)} & \varphi_{12}^{(3)} \\ \varphi_{21}^{(3)} & \varphi_{22}^{(3)} \\ 0 & 1 \end{pmatrix},$$

$$\begin{pmatrix} \varphi_{11}^{(4)} & \varphi_{12}^{(4)} \\ 1 & 0 \\ 0 & 1 \\ \varphi_{21}^{(4)} & \varphi_{22}^{(4)} \end{pmatrix}, \quad \begin{pmatrix} \varphi_{11}^{(5)} & \varphi_{12}^{(5)} \\ 1 & 0 \\ \varphi_{21}^{(5)} & \varphi_{22}^{(5)} \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} \varphi_{11}^{(6)} & \varphi_{12}^{(6)} \\ \varphi_{21}^{(6)} & \varphi_{22}^{(6)} \\ 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (3.13)$$

Together with the corresponding coordinates $\psi_{\alpha i}^{(A)}$ for the cotangent space in Eq. (2.13), these six sets of coordinate systems are sufficient to cover the entire manifold. Therefore this model has the six vacua given by

$$\langle \Phi \rangle = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (3.14)$$

which are the respective for the six choices given in (3.13), with $\langle \Psi \rangle = 0$. The set of two column vectors in each matrix in Eq. (3.14) constitutes a set of orthogonal basis vectors e_i chosen from the four basis vectors.

In the case $b \neq 0$, the superpotential (2.16) is

$$\begin{aligned} W &= b \sum_{a=1}^{N-1} \sum_{n=0}^{\infty} (-1)^n m_a \text{tr} \left[H_a \begin{pmatrix} (\psi\varphi)^n & (\psi\varphi)^n \psi \\ \varphi(\psi\varphi)^n & (\varphi\psi)^{n+1} \end{pmatrix} \right] \\ &= b \sum_{a=1}^{N-1} \sum_{n=0}^{\infty} (-1)^n m_a \text{tr} \left[H_a \begin{pmatrix} (\psi\varphi)^n & 0 \\ 0 & (\varphi\psi)^{n+1} \end{pmatrix} \right], \end{aligned} \quad (3.15)$$

where the last equality holds because the matrices H_a are diagonal. Similarly to the $T^*\mathbf{CP}^{N-1}$ case, the origin $\varphi = \psi^T = 0$ of each patch is a vacuum and we cannot have any other vacua.

§4. Massive HK quotient in the Wess-Zumino gauge

In the previous sections, we eliminated $\mathcal{N} = 2$ vector multiplets (as auxiliary superfields) without taking the Wess-Zumino gauge. In this section, we take the Wess-Zumino gauge, derive the bosonic action, and investigate vacua.

4.1. Lagrangian in the Wess-Zumino gauge

It is difficult to calculate the scalar potential without taking the Wess-Zumino gauge, because the inverse metric is difficult to obtain. On the other hand, in the Wess-Zumino gauge, the scalar potential can be directly obtained, while the Kähler potential and the superpotential cannot be calculated easily. We calculate the scalar

potential in this subsection. We represent the lowest components in the superfields using the same letters as we have used for the corresponding superfields.

In the Wess-Zumino gauge, the bosonic action of (2.3) is given by

$$\mathcal{L}_{\text{boson}} = \mathcal{L}_{\text{kin}} + \mathcal{L}_{\text{constr}} + \mathcal{L}_{\text{pot}}, \quad (4.1)$$

in which each term is given by

$$\begin{aligned} \mathcal{L}_{\text{kin}} &= -\text{tr} \left[(\partial_\mu \Phi^\dagger - i v_\mu \Phi^\dagger) (\partial^\mu \Phi + i \Phi v^\mu) \right] - \text{tr} \left[(\partial_\mu \Psi - i v_\mu \Psi) (\partial^\mu \Psi^\dagger + i \Psi^\dagger v^\mu) \right] \\ &= -\text{tr} (\partial_\mu \Phi^\dagger \partial^\mu \Phi + \partial_\mu \Psi \partial^\mu \Psi^\dagger) + i \text{tr} \left[v^\mu (\Phi^\dagger \overleftrightarrow{\partial}_\mu \Phi + \Psi \overleftrightarrow{\partial}_\mu \Psi^\dagger) \right] \\ &\quad - \text{tr} \left[v^\mu v_\mu (\Phi^\dagger \Phi + \Psi \Psi^\dagger) \right], \end{aligned} \quad (4.2)$$

$$\mathcal{L}_{\text{constr}} = \text{tr} [D(\Phi^\dagger \Phi - \Psi \Psi^\dagger - c \mathbf{1}_M)] + \text{tr} [F_\Sigma (\Psi \Phi - b \mathbf{1}_M) + \text{c.c.}], \quad (4.3)$$

$$\begin{aligned} \mathcal{L}_{\text{pot}} &= \text{tr} (F_\Phi^\dagger F_\Phi + F_\Psi^\dagger F_\Psi) \\ &\quad + \left[\text{tr} \{ \Sigma (F_\Psi \Phi + \Psi F_\Phi) \} + \sum_a m_a \text{tr} (F_\Psi H_a \Phi + \Psi H_a F_\Phi) + \text{c.c.} \right] \\ &\equiv -V(\Phi, \Psi, \Sigma), \end{aligned} \quad (4.4)$$

where we have defined $A \overleftrightarrow{\partial}_\mu B \equiv A(\partial_\mu B) - (\partial_\mu A)B$.

The equation of motion for the gauge field v_μ without the kinetic term in (4.1) reads

$$\partial \mathcal{L} / \partial v_\mu = -\{v_\mu, (\Phi^\dagger \Phi + \Psi \Psi^\dagger)\} + i(\Phi^\dagger \overleftrightarrow{\partial}_\mu \Phi + \Psi \overleftrightarrow{\partial}_\mu \Psi^\dagger) = 0. \quad (4.5)$$

Here $\{ , \}$ is an anti-commutator. This equation can be solved by expanding fields that takes values in the Lie algebra in terms of the $M \times M$ matrix T_A of the fundamental representation of $U(M)$ generators

$$\begin{aligned} v_\mu &\equiv v_\mu^A T_A, & \text{tr} (T_A T_B) &= \delta_{AB} \\ \Phi^\dagger \Phi + \Psi \Psi^\dagger &\equiv A^A T_A, & \tilde{A}_{AB} &\equiv \frac{1}{2} \text{tr} (\{T_A, T_B\} T_C) A^C, \\ i(\Phi^\dagger \overleftrightarrow{\partial}_\mu \Phi + \Psi \overleftrightarrow{\partial}_\mu \Psi^\dagger) &\equiv B_\mu^A T_A. \end{aligned} \quad (4.6)$$

Then we can express the gauge fields v_μ in terms of the dynamical scalar fields, solving the equation of motion (4.5) as

$$v_\mu^A = \frac{1}{2} \left(\tilde{A}^{-1} \right)_{AB} B_\mu^B. \quad (4.7)$$

If we eliminate the gauge field using this algebraic equation of motion, we obtain the kinetic term for hypermultiplets as

$$\mathcal{L}_{\text{kin}} = -\text{tr} (\partial_\mu \Phi^\dagger \partial^\mu \Phi + \partial_\mu \Psi \partial^\mu \Psi^\dagger) + \frac{1}{4} B^{A\mu} \left(\tilde{A}^{-1} \right)_{AB} B_\mu^B. \quad (4.8)$$

If we integrate the Lagrange multiplier fields D and F_Σ in (4.3), we obtain three real constraints

$$\Phi^\dagger \Phi - \Psi \Psi^\dagger = c \mathbf{1}_M, \quad \Psi \Phi = b \mathbf{1}_M. \quad (4.9)$$

The left-hand sides of these relations constitute the triplet of the moment map (Killing potential) for the $U(M)$ gauge symmetry. These values are fixed to the FI parameters, and the hyper-Kähler quotient is obtained together with the $U(M)$ quotient given in (4.7).

The Lagrangian (4.4) gives the following algebraic equations of motion for the auxiliary fields F_Φ and F_Ψ :

$$F_\Phi^\dagger = -\Sigma \Psi - \sum_a m_a \Psi H_a, \quad F_\Psi^\dagger = -\Phi \Sigma - \sum_a m_a H_a \Phi. \quad (4.10)$$

After eliminating the auxiliary fields F_Φ and F_Ψ using these algebraic equations of motion, we obtain the potential term V in Eq.(4.4) as

$$\begin{aligned} V(\Phi, \Psi, \Sigma) &= \text{tr} (F_\Phi^\dagger F_\Phi + F_\Psi^\dagger F_\Psi) \\ &= \text{tr} \left((\Sigma \Psi + \sum_a m_a \Psi H_a) (\Psi^\dagger \Sigma^\dagger + \sum_b m_b^* H_b \Psi^\dagger) \right. \\ &\quad \left. + (\Sigma^\dagger \Phi^\dagger + \sum_a m_a^* \Phi^\dagger H_a) (\Phi \Sigma + \sum_b m_b H_b \Phi) \right) \\ &= \text{tr} \left(\Sigma^\dagger \Sigma \Psi \Psi^\dagger + \sum_{ab} m_a m_b^* \Psi H_a H_b \Psi^\dagger \right) \\ &\quad + \text{tr} \left(\Sigma \Sigma^\dagger \Phi^\dagger \Phi + \sum_{ab} m_a^* m_b \Phi^\dagger H_a H_b \Phi \right) \\ &\quad + \sum_a \text{tr} \left((m_a^* \Sigma + m_a \Sigma^\dagger) (\Phi^\dagger H_a \Phi + \Psi H_a \Psi^\dagger) \right). \end{aligned} \quad (4.11)$$

Next, to eliminate Σ , we define the following quantities:

$$\begin{aligned} \Sigma &\equiv \Sigma^A T_A, & \Sigma^\dagger &\equiv \Sigma^{A*} T_A, \\ \Psi \Psi^\dagger &\equiv C^A T_A, & \tilde{C}_{AB} &\equiv \text{tr} (T_A T_B T_C) C^C, \end{aligned}$$

$$\begin{aligned}\Phi^\dagger\Phi &\equiv D^A T_A, & \tilde{D}_{AB} &\equiv \text{tr} (T_B T_A T_C) D^C, \\ \sum_a m_a^* \left(\Phi^\dagger H_a \Phi + \Psi H_a \Psi^\dagger \right) &\equiv E^A T_A.\end{aligned}\quad (4.12)$$

Because the equation of motion for Σ is purely algebraic, we can eliminate it as

$$\Sigma^A = - \left((\tilde{C} + \tilde{D})^{-1} \right)_{AB} E^{B*}, \quad \Sigma^{A*} = -E^B \left((\tilde{C} + \tilde{D})^{-1} \right)_{BA}. \quad (4.13)$$

After eliminating the auxiliary fields Σ in the $\mathcal{N} = 2$ vector multiplet (V, Σ) using its algebraic equation of motion, we obtain the potential in terms of dynamical scalar fields,

$$\begin{aligned}V(\Phi, \Psi) &= \text{tr} \left(\sum_{ab} \left(m_a m_b^* \Psi H_a H_b \Psi^\dagger + m_a^* m_b \Phi^\dagger H_a H_b \Phi \right) \right) \\ &\quad - E^A \left((\tilde{C} + \tilde{D})^{-1} \right)_{AB} E^{B*}.\end{aligned}\quad (4.14)$$

4.2. Vacua in the massive $T^*\mathbf{CP}^{N-1}$ model

We first focus on the vacua of the $T^*\mathbf{CP}^{N-1}$ model (the $M=1$ case). Here, we are interested in the SUSY vacua. SUSY vacua correspond to vanishing auxiliary fields F_Φ and F_Ψ , i.e.,

$$0 = -F_{\Phi^i}^\dagger = \Psi_i (\Sigma + \tilde{m}_i), \quad (4.15)$$

$$0 = -F_{\Psi_i}^\dagger = (\Sigma + \tilde{m}_i) \Phi^i, \quad (4.16)$$

and the constraints

$$0 = \Phi_i^\dagger \Phi^i - \Psi_i \Psi^{\dagger i} - c, \quad (4.17)$$

$$0 = \Psi_i \Phi^i - b, \quad (4.18)$$

where we have defined $\tilde{m}_i$ by $\sum_a m_a H_a = \text{diag.}(\tilde{m}_1, \tilde{m}_2, \dots, \tilde{m}_N)$, with $\sum_{i=1}^N \tilde{m}_i = 0$. We assume a generic mass $\tilde{m}_i \neq \tilde{m}_j$ ^{††} and that either b or c has a nonzero value.

^{††} This assumption is the same as the assumption $M_a \neq 0$, which was made in the study of the vacua in §3.1. Indeed, the condition $\tilde{m}_i \neq \tilde{m}_j$ for any combination of i and j ($i \neq j$) is equivalent to the condition $M_a \neq 0$ for all a in all patches. If $\tilde{m}_i = \tilde{m}_j$ holds for a pair i and j , there is a patch where $M_a = 0$ for one a . The vacuum is then not localized at the origin of the patch, and hence continuous vacua appear. Therefore, the discussion in §3.1 no longer holds. For instance, in the $N = 3$ case, the relation $M_a = \tilde{m}_3 - \tilde{m}_a$ holds in the patch, and therefore we have $\Phi^t = (\varphi, 1)^t$ and $\Psi = (\psi, -\psi\varphi)$ instead of (2.13). It is easily seen that $M_2 = 0$ if $\tilde{m}_2 = \tilde{m}_3$, while $M_1 \neq 0$.

Equation (4.15) implies $\Psi_i = 0$ or

$$\Sigma = -\tilde{m}_i, \quad (4.19)$$

with $\Psi_i \neq 0$ for some i ($= 1, 2, \dots, N$). In the case $\Sigma \neq -\tilde{m}_i$ for all i , we have $\Phi = 0$ and $\Psi = 0$, from Eqs. (4.15) and (4.16). However, $\Phi = \Psi^T = 0$ is inconsistent with Eq. (4.17) for $c \neq 0$ or with Eq. (4.18) for $b \neq 0$. Therefore Eq. (4.19) must hold for some i , and Φ^i and/or Ψ_i can be nonzero for such i . Hence, there exist N vacua labelled by $i = 1, \dots, N$. To determine the vacuum expectation values of Φ^i and Ψ_i for the i -vacuum, let us assume $b = 0$ and $c \neq 0$, without loss of generality. Then, from Eq. (4.18) with $b = 0$, Φ^i or Ψ_i must be zero. Using Eq. (4.17), we obtain $|\Phi^i| = \sqrt{c}$ and $\Psi_i = 0$ if $c > 0$ or $\Phi^i = 0$, and $|\Psi_i| = \sqrt{-c}$ if $c < 0$. In the case $c = 0$ and $b \neq 0$, we obtain $\Phi^i = \sqrt{b}e^{i\theta}$ and $\Psi_i = \sqrt{b}e^{-i\theta}$ with an arbitrary phase θ . We thus have found N discrete vacua.

4.3. Vacua in the massive $T^*G_{N,M}$ model

The analysis of vacua for the massive $T^*\mathbf{CP}^{N-1}$ model in the previous subsection can be generalized to the case of the massive $T^*G_{N,M}$ model. The vacuum conditions in this case are given by

$$0_{\alpha i} = -F_{\Phi\alpha i}^\dagger = (\Sigma_\alpha^\beta + \delta_\alpha^\beta \tilde{m}_i)\Psi_{\beta i}, \quad (4.20)$$

$$0^{i\alpha} = -F_\Psi^{\dagger i\alpha} = \Phi^{i\beta}(\Sigma_\beta^\alpha + \delta_\beta^\alpha \tilde{m}_i), \quad (4.21)$$

along with the constraints

$$0_\alpha^\beta = \Phi_{\alpha i}^\dagger \Phi^{i\beta} - \Psi_{\alpha i} \Psi^{\dagger i\beta} - c\delta_\alpha^\beta, \quad (4.22)$$

$$0_\alpha^\beta = \Psi_{\alpha i} \Phi^{i\beta} - b\delta_\alpha^\beta, \quad (4.23)$$

where $\alpha, \beta = 1, \dots, M$ are gauge indices. Using the $U(M)$ gauge rotation, Σ can be diagonalized:

$$\Sigma = \text{diag.}(\Sigma_1, \Sigma_2, \dots, \Sigma_M), \quad (4.24)$$

$$\Sigma_1 + \Sigma_2 + \dots + \Sigma_M = M\Sigma^0. \quad (4.25)$$

To infer the consequences of Eqs. (4.20) and (4.21), we should consider two cases separately, that in which $\Sigma_\alpha + \tilde{m}_i \neq 0$ for some α and all i and that in which

$\Sigma_\alpha + \tilde{m}_i = 0$ for all α and some i . In the former case, we get $\Phi = 0$ and $\Psi = 0$. However, these are inconsistent with (4.22) for $c \neq 0$ and with (4.23) for $b \neq 0$, and hence they do not represent a solution. In the latter case, let us assume the generic case for masses, i.e. $\tilde{m}_i \neq \tilde{m}_j$.^{†††} The eigenvalue equation (4.21) for Σ_α gives the eigenvalue $\Sigma_\alpha = -\tilde{m}_i$ corresponding to the eigenvector formed by the α -th column vector, $(\Phi^{1\alpha}, \dots, \Phi^{N\alpha})^T$, with only a single nonvanishing element. Similarly, the eigenvalue equation (4.20) gives the same eigenvalue $\Sigma_\alpha = -\tilde{m}_i$ corresponding to the eigenvector formed by the α -th row vector, $(\Psi_{\alpha 1}, \dots, \Psi_{\alpha N})$, with only a single nonvanishing element. Explicitly, we have

$$\Phi = \begin{pmatrix} 0 \\ \vdots \\ \Phi^{i\alpha} \\ 0 \\ \vdots \end{pmatrix}, \quad \Psi = \begin{pmatrix} 0 & \dots & \Psi_{\alpha i} & 0 & \dots \end{pmatrix}. \quad (4.26)$$

Since there are N different masses $\tilde{m}_i$, there are N corresponding nontrivial eigenvectors of the form given in (4.26). The diagonal Σ_α should be equal to one such mass, $-\tilde{m}_i$. We have M different diagonal elements Σ_α ($\alpha = 1, \dots, M$). Therefore, we obtain ${}_NC_M$ possibilities of different vacua, counting a vacua related by relabelling of the gauge indices α as equivalent. The values of the nonvanishing elements $\Phi^{i\alpha}$ and $\Psi_{\alpha i}$ of the eigenvectors in Eq. (4.26) are explicitly determined by the constraints (4.22) and (4.23):

$$|\Phi^{i\alpha}|^2 - |\Psi_{\alpha i}|^2 = c, \quad (4.27)$$

$$\Psi_{\alpha i} \Phi^{i\alpha} = b \text{ (no sum)}. \quad (4.28)$$

Let us again assume $b = 0$ and $c \neq 0$. Then, if $c > 0$, we have $|\Phi^{i\alpha}| = \sqrt{c}$ and $\Psi_{\alpha i} = 0$, and if $c < 0$, we have $\Phi^{i\alpha} = 0$ and $|\Psi_{\alpha i}| = \sqrt{-c}$. Thus, we obtain the same result as in the previous section. In the case $b \neq 0$ and $c = 0$, we have $\Phi^{i\alpha} = \sqrt{b}e^{i\theta}$ and $\Psi_{\alpha i} = \sqrt{b}e^{-i\theta}$, with an arbitrary phase θ .

For example, let us consider the case $M = 2$, $N = 4$ again. In this case, there

^{†††} As in the case of the $T^*\mathbf{CP}^{N-1}$ model, it can be confirmed that this assumption is the same as $M_{ai} \neq 0$ used in §3.2.

are ${}_4C_2 = 6$ solutions for both Φ and Ψ :

$$\begin{aligned} \Phi = & \begin{pmatrix} \Phi^{11} & 0 \\ 0 & \Phi^{22} \\ 0 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} \Phi^{11} & 0 \\ 0 & 0 \\ 0 & \Phi^{32} \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} \Phi^{11} & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & \Phi^{42} \end{pmatrix}, \\ & \begin{pmatrix} 0 & 0 \\ \Phi^{21} & 0 \\ 0 & \Phi^{32} \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ \Phi^{21} & 0 \\ 0 & 0 \\ 0 & \Phi^{42} \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ \Phi^{31} & 0 \\ 0 & \Phi^{42} \end{pmatrix}, \end{aligned} \quad (4.29)$$

and

$$\begin{aligned} \Psi = & \begin{pmatrix} \Psi_{11} & 0 & 0 & 0 \\ 0 & \Psi_{22} & 0 & 0 \end{pmatrix}, \begin{pmatrix} \Psi_{11} & 0 & 0 & 0 \\ 0 & 0 & \Psi_{23} & 0 \end{pmatrix}, \begin{pmatrix} \Psi_{11} & 0 & 0 & 0 \\ 0 & 0 & 0 & \Psi_{24} \end{pmatrix}, \\ & \begin{pmatrix} 0 & \Psi_{12} & 0 & 0 \\ 0 & 0 & \Psi_{23} & 0 \end{pmatrix}, \begin{pmatrix} 0 & \Psi_{12} & 0 & 0 \\ 0 & 0 & 0 & \Psi_{24} \end{pmatrix}, \begin{pmatrix} 0 & 0 & \Psi_{13} & 0 \\ 0 & 0 & 0 & \Psi_{24} \end{pmatrix}. \end{aligned} \quad (4.30)$$

Clearly, for $b = 0$ and $c \neq 0$, we obtain $|\Phi^{i\alpha}| = \sqrt{c}$ and $\Psi_{\alpha i} = 0$ if $c > 0$ and $\Phi^{i\alpha} = 0$ and $|\Psi_{\alpha i}| = \sqrt{-c}$ if $c < 0$.

§5. Massive HK quotient with respect to $SU(M)$ gauge group

In this subsection, we construct the massive HK sigma model with the $SU(M)$ gauge group. We eliminate the vector multiplets in the superfield formalism and find that this model does not have discrete vacua. Because we can carry out the same analysis in the case of Wess-Zumino gauge as in the case of the $U(M)$ gauge group, we do not repeat it.

5.1. Massive HK Sigma Model with SU gauge group

In this section, we consider $\mathcal{N} = 2$ SUSY QCD with N flavors and the $SU(M)$ gauge group. We employ the same matter field content as with $T^*G_{N,M}$, but here, the gauge multiplets take values in the Lie algebra $SU(M)$, i.e., $V = V^A T_A$ and $\Sigma = \Sigma^A T_A$, where T_A represents the generators of $SU(M)$. Then, the Lagrangian

is given by

$$\begin{aligned} \mathcal{L} = & \int d^4\theta \left[\text{tr}(\Phi^\dagger \Phi e^V) + \text{tr}(\Psi \Psi^\dagger e^{-V}) \right] \\ & + \left[\int d^2\theta \left(\text{tr}(\Sigma \Psi \Phi) + \sum_{a=1}^{N-1} m_a \text{tr}(\Psi H_a \Phi) \right) + \text{c.c.} \right]. \end{aligned} \quad (5.1)$$

There are no FI parameters, because there is no $U(1)$ gauge symmetry. The $SU(M)$ gauge transformation is obtained in the same way as in the $U(M)$ case, and it is complexified to $SU(M)^{\mathbf{C}} = SL(M, \mathbf{C})$. This model has the additional $U(1)_{\text{D}}$ flavor symmetry

$$\Phi \rightarrow \Phi' = e^{i\lambda} \Phi, \quad \Psi \rightarrow \Psi' = e^{-i\lambda} \Psi, \quad (5.2)$$

which was gauged in the $U(M)$ case. In the massless limit, $m_a \rightarrow 0$, the total flavor symmetry is $U(N) = SU(N) \times U(1)_{\text{D}}$. With a non-vanishing mass term, this $U(N)$ is explicitly broken down to $U(1)^N$.

We eliminate all auxiliary superfields in the superfield formalism. Then, the equations of motion for V and Σ read^x

$$\frac{\partial \mathcal{L}}{\partial X^A} = \text{tr}[(\Phi^\dagger \Phi e^V - e^{-V} \Psi \Psi^\dagger) T_A] = 0, \quad (5.3)$$

$$\frac{\partial \mathcal{L}}{\partial \Sigma^A} = \text{tr}(\Psi \Phi T_A) = 0, \quad (5.4)$$

respectively. These equations imply

$$\Phi^\dagger \Phi e^V - e^{-V} \Psi \Psi^\dagger = C \mathbf{1}_M, \quad (5.5)$$

$$\Psi \Phi = B \mathbf{1}_M, \quad (5.6)$$

respectively, where $C(x, \theta, \bar{\theta})$ and $B(x, \theta, \bar{\theta})$ are vector and chiral superfields in the $\mathcal{N} = 1$ superfield formalism.^{xi}

The gauge field V can be obtained in terms of the dynamical fields from Eq. (5.5) as

$$e^V = \frac{1}{2} (\Phi^\dagger \Phi)^{-1} \left(C \mathbf{1}_M \pm \sqrt{C^2 \mathbf{1}_M + 4 \Phi^\dagger \Phi \Psi \Psi^\dagger} \right). \quad (5.7)$$

^x Here we take the Maurer-Cartan 1-form $\delta X = \delta X^A T_A \equiv e^{-V} \delta e^V$ as an infinitesimal parameter of variation, as in the case of $T^* G_{N,M}$. Then, the equations $\delta \text{tr}(\Phi^\dagger \Phi e^V) = \text{tr}(\Phi^\dagger \Phi e^V T_A) \delta X^A$ and $\delta \text{tr}(\Psi \Psi^\dagger e^{-V}) = -\text{tr}(e^{-V} \Psi \Psi^\dagger T_A) \delta X^A$ hold.

^{xi} We can understand how C satisfies the condition for a vector superfield, $C^\dagger = C$, if we rewrite Eq. (5.5) as $e^{\frac{V}{2}} \Phi^\dagger \Phi e^{\frac{V}{2}} - e^{-\frac{V}{2}} \Psi \Psi^\dagger e^{-\frac{V}{2}} = C \mathbf{1}_M$.

Then, because the equation $\det e^V = 1$ holds, we obtain the equation

$$\det \left(C \mathbf{1}_M \pm \sqrt{C^2 \mathbf{1}_M + 4\Phi^\dagger \Phi \Psi \Psi^\dagger} \right) = 2^M \det(\Phi^\dagger \Phi), \quad (5.8)$$

which enables us to express C in terms of dynamical fields implicitly: $C = C(\Phi, \Phi^\dagger; \Psi, \Psi^\dagger)$.

On the other hand, Eq. (5.6) implies

$$B = \frac{1}{M} \text{tr}(\Phi \Psi). \quad (5.9)$$

Substituting the solution (5.7) back into the Lagrangian (5.1), we obtain the Kähler potential

$$K = \pm \text{tr} \sqrt{C^2(\Phi, \Phi^\dagger; \Psi, \Psi^\dagger) \mathbf{1}_M + 4\Phi^\dagger \Phi \Psi \Psi^\dagger}, \quad (5.10)$$

with C satisfying the constraint (5.8). We should choose the plus sign to realize the positivity of the metric. Using the $N \times N$ meson matrix $\mathcal{M} = \Phi \Psi$, the Kähler potential can be rewritten as⁸⁾

$$K = \frac{M}{N} \text{tr}_{N \times N} \sqrt{C^2 \mathbf{1}_N + 4 \frac{N}{M} \mathcal{M} \mathcal{M}^\dagger}. \quad (5.11)$$

The Kähler potential (5.10) or (5.11) is strictly invariant under the full global (flavor) $U(N)$ symmetry, because the mass term in the Lagrangian (5.1) does not affect the D-term.

Let us stipulate the complex gauge symmetry $SU(M)^{\mathbf{C}} = SL(M, \mathbf{C})$ to express the Lagrangian in terms of independent superfields. We can employ a gauge that is similar to that in the $b \neq 0$ case in $T^*G_{N,M}$:

$$\Phi = \sigma \begin{pmatrix} \mathbf{1}_M \\ \varphi \end{pmatrix} P, \quad \Psi = P(\mathbf{1}_M, \psi) \rho, \quad P = (\mathbf{1}_M + \psi \varphi)^{-\frac{1}{2}}. \quad (5.12)$$

Here φ and ψ are $[(N - M) \times M]$ - and $[M \times (N - M)]$ -matrix chiral superfields, respectively, and σ and ρ are chiral superfields satisfying $\sigma \rho = B$ from Eq. (5.9). We can consider σ and ρ to be independent fields among the three fields σ, ρ and B .

Substituting Eq. (5.12) into the Kähler potential (5.10), we obtain the Kähler potential in terms of the independent fields $\varphi, \psi, \rho, \sigma$ and their conjugates. The superpotential also can be calculated as

$$W = \sum_a m_a \sigma \rho \text{tr} \left[H_a \begin{pmatrix} \mathbf{1}_M \\ \varphi \end{pmatrix} (\mathbf{1}_M + \psi \varphi)^{-1} (\mathbf{1}_M, \psi) \right]. \quad (5.13)$$

This target manifold has the isometry $U(N) = SU(N) \times U(1)_D$, in which the $SU(N)$ part is the same as that in the case of $T^*G_{N,M}$. The Kähler potential does not undergo the Kähler transformation. As for the symmetry of the Lagrangian, the superpotential is invariant under the $U(1)$ fiber symmetry originating from (5.2),

$$\sigma \rightarrow \sigma' = e^{i\lambda} \sigma, \quad \rho \rightarrow \rho' = e^{-i\lambda} \rho, \quad (5.14)$$

in addition to the $U(1)^{N-1}$ symmetry of the massive $T^*G_{N,M}$ model. Gauging this $U(1)_D$ symmetry, we obtain the $T^*G_{N,M}$ model again, by definition. Gauging the $U(1)_D$ symmetry implies fixing B and C in the constraints (5.8) and (5.9) as constants. Then, these constraints fixing become identical to those in the $T^*G_{N,M}$ case (2.5) and (2.6), respectively.

The above discussion clarifies the bundle structure: The set of σ and ρ constitutes a fiber of quaternion with manifold as a whole being the quaternionic line bundle over $T^*G_{N,M}$.

It is interesting that we can define the same model using a Lagrangian similar to that in the $T^*G_{N,M}$ case if we promote the FI-parameters b and c in the $T^*G_{N,M}$ model to the chiral and vector superfields $B(x, \theta, \bar{\theta})$ and $C(x, \theta, \bar{\theta})$, respectively, constituting an $\mathcal{N} = 2$ vector multiplet without a kinetic term. Then, the Lagrangian has precisely the same field content as that for $T^*G_{N,M}$ ^{xii}:

$$\begin{aligned} \mathcal{L} = & \int d^4\theta \left[\text{tr}(\Phi^\dagger \Phi e^V) + \text{tr}(\Psi \Psi^\dagger e^{-V}) - C \text{tr} V \right] \\ & + \left[\int d^2\theta \left(\text{tr} \{ \Sigma(\Psi \Phi - B \mathbf{1}_M) \} + \sum_{a=1}^{N-1} m_a \text{tr}(\Psi H_a \Phi) \right) + \text{c.c.} \right]. \end{aligned} \quad (5.15)$$

In this case, the equations of motion for V , Σ , B and C read

$$\frac{\partial \mathcal{L}}{\partial V} = \Phi^\dagger \Phi e^V - e^{-V} \Psi \Psi^\dagger - C \mathbf{1}_M = 0, \quad \frac{\partial \mathcal{L}}{\partial \Sigma} = \Psi \Phi - B \mathbf{1}_M = 0, \quad (5.16)$$

$$\frac{\partial \mathcal{L}}{\partial B} = -\text{tr} \Sigma = 0, \quad \frac{\partial \mathcal{L}}{\partial C} = -\text{tr} V = 0. \quad (5.17)$$

We thus have the same constraints as in the model (5.1). The technique of promoting an FI-parameter to a superfield was used in Ref. 44) to construct the complex

^{xii} $\mathcal{N} = 2$ SUSY with this action can be understood by considering the transformation of the $U(1)_J$ group which is the subgroup of $SU(2)_R$ under which $\theta \rightarrow e^{i\alpha} \theta$, $\Phi \rightarrow e^{i\alpha} \Phi$, $\Psi \rightarrow e^{-i\alpha} \Psi$, $V \rightarrow V$, $\Sigma \rightarrow \Sigma$, $C \rightarrow C$ and $B \rightarrow B$.

canonical line bundle over $G_{N,M}$ in an $\mathcal{N} = 1$ (four SUSY) theory, where an additional superfield σ relative to $G_{N,M}$ appeared and was identified with the fiber of a complex line. Returning to $\mathcal{N} = 2$, we obtain the total manifold as the cotangent bundle over the complex line bundle on $G_{N,M}$, where $\rho(\psi)$ serves as the cotangent fiber over $\sigma(\varphi)$. We have thus found that the space possesses two equivalent bundle structures. We may need to transform σ and ρ to some fields σ' and ρ' in order to avoid coordinate singularities, as in the four SUSY case.⁴⁴⁾

Other gauge groups: SO and Sp

The equations of motion for V and Σ takes the same form as Eqs. (5.3) and (5.4) for *any* gauge group. However, the right-hand side of Eq.(5.5) must be a symmetric tensor for the $SO(M)$ gauge group, and the right-hand side of Eq.(5.6) must be a pseudo-symmetric tensor for the $Sp(M)$ gauge group.

5.2. Vacua of SU gauge theories

We now seek the vacua of the HK sigma model with the SU gauge group. The superpotential (5.13) of this model can be rewritten as

$$W = \sigma\rho \sum_{a=1}^{N-1} \sum_{n=0}^{\infty} (-1)^n m_a \text{tr} \left[H_a \begin{pmatrix} (\psi\varphi)^n & 0 \\ 0 & (\varphi\psi)^{n+1} \end{pmatrix} \right] \equiv \sigma\rho W_U, \quad (5.18)$$

where W_U (times b) denotes the superpotential (2.16) or (3.15) of the $U(M)$ gauge group with $b \neq 0$. The derivatives of the superpotential with respect to the fields are given by $\partial_\psi W = \sigma\rho\partial_\psi W_U$, $\partial_\varphi W = \sigma\rho\partial_\varphi W_U$, $\partial_\rho W = \sigma W_U$ and $\partial_\sigma W = \rho W_U$. The vacuum condition is given by $\sigma = \rho = 0$, because $\partial W_U = 0$ holds only at $\varphi = \psi^T = 0$, as found in the previous section, but we have $W_U \neq 0$ there. Therefore, this model has no discrete vacua, and so there cannot be any wall solution.

Vacua of the massive HK sigma model with the $SU(M)$ gauge group may have a similarity with that with the $U(M)$ gauge group, the massive $T^*G_{N,M}$ model, in a particular limit. In the latter model, there exist the FI-parameters b and c , which represent the radius of the base $G_{N,M}$, and the discrete vacua are attached to this manifold. In the limit of vanishing b and c , a singularity appears. All of the discrete vacua go to that point, and as a result there are no sets of discrete vacua. Because the massive HK sigma model with the $SU(M)$ gauge group cannot contain the FI parameters from the beginning, we believe that this model has no discrete vacua. It

is thus seen that the FI-parameters seem to play an essential role in determining the existence of discrete vacua.

§6. Formulation in the harmonic superspace

In this section we formulate our models with $U(M)$ and $SU(M)$ gauge symmetries using the HSF. Massless nonlinear sigma models with toric and non-toric HK manifolds using the HSF were originally considered in Refs. 46) and 47) using the quotient construction. Massive nonlinear sigma models with toric HK manifolds, like the Taub-NUT⁴⁸⁾ metric and the Eguchi-Hanson metric, have also been studied.^{11),48)} We construct massive HK sigma models with our non-toric HK manifolds using HSF. One advantage of constructing massive nonlinear sigma models using the HSF is that $\mathcal{N} = 2$ SUSY is manifest in this case. It is easy to extend these models to cases with other flavor and gauge symmetries.

We first present the actions of the $U(M)$ and $SU(M)$ models in terms of the harmonic superfields. Next, we derive their component actions in the Wess-Zumino gauge. We find that the scalar potential is represented by auxiliary fields in the hypermultiplet analytic superfield and we obtain a vacuum condition that is identical to that in the $\mathcal{N} = 1$ formulation. Finally, we obtain the same result regarding the vacua as in the $\mathcal{N} = 1$ formulation. We follow the notation of Ref. 11).

6.1. Massive HK sigma model with $U(M)$ and $SU(M)$ gauge groups

First of all, we consider the model with $U(M)$ gauge symmetry. The HSF action for the sigma models with $U(M)$ and $SU(M)$ gauge symmetries can be described by two kinds of analytic superfields, which are hypermultiplet and vector multiplet analytic superfields. A hypermultiplet is defined by the analytic superfield $q^{+a\alpha}$ ($a = 1, \dots, N$, $\alpha = 1, \dots, M$) in the fundamental representation of the $SU(N)$ flavor symmetry^{xiii} and the $U(M)$ gauge symmetry, which is a function in the harmonic analytic $\mathcal{N} = 2$ superspace

$$\zeta_A = (x_A^\mu, \theta^+, \bar{\theta}^+, u_i^\pm), \quad (6.1)$$

^{xiii} To avoid confusion with the $SU(2)_R$ indices $i = 1, 2$ of the harmonic variable, we denote the $SU(N)$ flavor indices by $a = 1, \dots, N$ in the following.

where the coordinates $u^{+i}, u^{-i}, u^{+i}u_i^- = 1$ ($i = 1, 2$) are the $SU(2)_R/U(1)_r$ harmonic variables. Here $U(1)_r$ is a diagonal subgroup of $SU(2)_R$. The $U(M)$ gauge transformation for the hypermultiplet is given by

$$q^{+a\alpha} \rightarrow (e^{-i\lambda(\zeta_A, u)})^\alpha_\beta q^{+a\beta}, \quad (6.2)$$

where $\lambda = \lambda^A T_A$ [T_A being the generator of the $U(M)$ gauge symmetry] is the real analytic superfield with $U(1)_r$ charge 0, which represents the gauge transformation parameter. The vector multiplet $V^{++} = V^{A++}T_A$ is defined as the real analytic superfield, that is, $\widetilde{V^{++}} = V^{++}$, where the tilde denotes conjugation which is the combination of complex conjugation and the star conjugation.^{(11), (37)} The action of the tilde conjugation on ζ_A is defined as

$$\widetilde{x_A^\mu} = x_A^\mu, \quad \widetilde{\theta^+} = \bar{\theta}^+, \quad \widetilde{\bar{\theta}^+} = -\theta^+, \quad \widetilde{u_i^\pm} = u^{\pm i}, \quad \widetilde{u^{\pm i}} = u_i^\pm. \quad (6.3)$$

The vector multiplet superfield transforms under the $U(M)$ gauge transformation as

$$V^{++} \rightarrow e^{-i\lambda} V^{++} e^{i\lambda} - i e^{-i\lambda} D^{++} e^{i\lambda}, \quad (6.4)$$

where D^{++} is the covariant derivative defined by

$$D^{++} = \partial^{++} - 2i\theta^+ \sigma^\mu \bar{\theta}^+ \partial_\mu^A, \quad \partial^{++} = u_i^+ \frac{\partial}{\partial u_i^-}, \quad \partial_\mu^A = \frac{\partial}{\partial x_A^\mu}. \quad (6.5)$$

The infinitesimal forms of (6.2) and (6.4) are

$$\delta q^+ = -i\lambda q^+, \quad (6.6)$$

$$\delta V^{++} = D^{++} \lambda - i[\lambda, V^{++}]. \quad (6.7)$$

The action of the massive $T^*G_{N,M}$ model is invariant under these transformations, and it is given by

$$S = - \int d\zeta_A^{(-4)} du \left\{ \widetilde{q^+}_{a\alpha} (D^{++} + iV^{++})^\alpha_\beta q^{+a\beta} + \xi^{++} \text{tr}(V^{++}) - \sum_{A=1}^{N-1} \widetilde{q^+}_{a\alpha} (\theta^{+2} \bar{m}_A - \bar{\theta}^{+2} m_A) (H_A)^a_b q^{+b\alpha} \right\}, \quad (6.8)$$

where $d\zeta_A^{(-4)} du = d^4 x_A d^2 \theta^+ d^2 \bar{\theta}^+ du$ is the measure of the integration over analytic superspace given (6.1), $\xi^{++} = \xi^{(ij)} u_i^+ u_j^+$ is the coefficient of the FI term [which is

the $SU(2)_R$ triplet], m_A represents complex mass parameters, and H_A represents the diagonal generators of the Cartan subalgebra of $SU(N)$ defined in (2.3). Here, the tilde conjugation acts not on the gauge and flavor indices but on the $U(1)_r$ charge. The quantity V^{++} serves as the Lagrange multiplier. Integrating the vector multiplet, it gives a constraint for the hypermultiplet. At the component level, it gives a constraint on the scalars and fermions, and it makes the target manifold nontrivial, as we see in the next subsection. We have absorbed a common hypermultiplet mass through a shift of the analytic superfield V^{++} . In the limit $m_A \rightarrow 0$, the action (6.8) becomes that of the massless $T^*G_{N,M}$ model, whose isometry is $SU(N)$. The mass term explicitly breaks $SU(N)$ down to $U(1)^{N-1}$. This feature is identical to that in the case of the $\mathcal{N} = 1$ formalism.

Next we consider the model with $SU(M)$ gauge symmetry. This quotient action can be easily obtained by restricting the gauge symmetry $U(M)$ to $SU(M)$ in the action (6.8). In this case, the FI term does not exist, since the theory no longer possesses $U(1)$ gauge symmetry. The action is given by

$$S = - \int d\zeta_A^{(-4)} du \{ \widetilde{q}^+_{a\alpha} (D^{++} + iV^{++})^{\alpha}_{\beta} q^{+a\beta} - \sum_{A=1}^{N-1} \widetilde{q}^+_{a\alpha} (\theta^{+2} \bar{m}_A - \bar{\theta}^{+2} m_A) (H_A)^a_b q^{+b\alpha} \}. \quad (6.9)$$

Note again that the theory has an additional $U(1)$ global symmetry, like the $\mathcal{N} = 1$ formulation. The total global symmetry in the $m_A = 0$ case is $U(N)$. The global symmetry $U(N)$ is broken down to $U(1)^N$ when there exists a non-vanishing mass term.

We can obtain the nonlinear sigma model action in terms of the independent harmonic superfields by fixing the gauge symmetry and solving the constraint, as in the $\mathcal{N} = 1$ formulation. The resulting action can be described by the independent analytic superfields of hypermultiplets. However, it is often difficult to decompose the action consisting independent superfields into a component action, because the kinematical part of the equations of motion, which is necessary to obtain the component action, is difficult to solve. Instead, we can easily solve the kinematical part of the equations of motion when we employ the Wess-Zumino gauge and can obtain the explicit form of the component action. This gauge is particularly convenient to

understand the vacua of theories in the HSF.

6.2. Scalar potential in the Wess-Zumino gauge

We now focus on the $U(M)$ case. In the following, we write the mass matrix as $\Sigma_{A=1}^{N-1} m_A H_A \equiv \tilde{m}$ and $\Sigma_{A=1}^{N-1} \bar{m}_A H_A \equiv \tilde{\bar{m}}$. In this case, the action can be rewritten as

$$S = \int d\zeta_A^{(-4)} du \left\{ \bar{q}_{a\alpha}^+ (D_{cc}^{++} + iV^{++})^\alpha_{\beta} q^{+a\beta} - \xi^{++} \text{tr}(V^{++}) \right\}, \quad (6.10)$$

where we have used $\widetilde{q^+} = -\bar{q}^+$ and the definition of the covariant derivative including the mass term

$$D_{cc}^{++} \equiv \partial^{++} - 2i\theta^+ \sigma^\mu \bar{\theta}^+ \partial_\mu^A - (\theta^{+2} \tilde{\bar{m}} - \bar{\theta}^{+2} \tilde{m}). \quad (6.11)$$

The equations of motion are

$$0 = (D_{cc}^{++} + iV^{++})^\alpha_{\beta} q^{+a\beta}, \quad (6.12)$$

$$0 = \bar{q}_{a\alpha}^+ i(T_A)^\alpha_{\beta} q^{+a\beta} - \xi^{++} \text{tr}(T_A), \quad (6.13)$$

where (6.12) includes kinematical and dynamical parts of the equations of motion, and (6.13) is a constraint. In the HK sigma model, we are only interested in the bosonic components. To obtain the bosonic Lagrangian, we have to solve the kinematical part of the equations of motion to eliminate the infinite set of auxiliary fields. To solve the kinematical part, we substitute the Grassmann expansion of the analytic superfields into the above equations. In the Wess-Zumino gauge, the bosonic components in the Grassmann expansion of the analytic superfields $q^{+a\alpha}$ and V^{++} are given by

$$q^{+a\alpha} = F^{+a\alpha} + \theta^{+2} M^{-a\alpha} + \bar{\theta}^{+2} N^{-a\alpha} + i\theta^+ \sigma^\mu \bar{\theta}^+ A_\mu^{-a\alpha} + \theta^{+2} \bar{\theta}^{+2} D^{(-3)a\alpha}, \quad (6.14)$$

$$V^{++} = \theta^{+2} \bar{M}_v + \bar{\theta}^{+2} M_v + i\theta^+ \sigma^\mu \bar{\theta}^+ (-2V_\mu) + \theta^{+2} \bar{\theta}^{+2} D_v^{(--)}, \quad (6.15)$$

where each component in V^{++} is a Lie algebra-valued field, for example $M_v = M_v^A T_A$, and $D_v^{(--)} \equiv D_v^{(ij)} u_i^- u_j^-$. Note that the components in V^{++} do not depend on the harmonic variables, whereas the components in q^+ include the harmonic variables. Substituting them into (6.12), we obtain the following kinematical part of

the equations of motion

$$0 = \partial^{++} F^{+a\alpha}, \quad (6.16)$$

$$0 = \partial^{++} M^{-a\alpha} - \tilde{m}^a_b F^{+b\alpha} + i\bar{M}_v^\alpha{}_\beta F^{+a\beta}, \quad (6.17)$$

$$0 = \partial^{++} N^{-a\alpha} + \tilde{m}^a_b F^{+b\alpha} + iM_v^\alpha{}_\beta F^{+a\beta}, \quad (6.18)$$

$$0 = \partial^{++} A_\mu^{-a\alpha} - 2\partial_\mu^A F^{+a\alpha} - 2iV_\mu^\alpha{}_\beta F^{+a\beta}. \quad (6.19)$$

The equations (6.16)–(6.19) can be easily solved, and we find

$$F^{+a\alpha}(x_A, u) = f^{ia\alpha}(x_A)u_i^+, \quad (6.20)$$

$$M^{-a\alpha}(x_A, u) = \tilde{m}^a_b f^{ib\alpha}(x_A)u_i^- - i(\bar{M}_v)^\alpha{}_\beta f^{ia\beta}(x_A)u_i^-, \quad (6.21)$$

$$N^{-a\alpha}(x_A, u) = -\tilde{m}^a_b f^{ib\alpha}(x_A)u_i^- - i(M_v)^\alpha{}_\beta f^{ia\beta}(x_A)u_i^-, \quad (6.22)$$

$$A_\mu^{-a\alpha}(x_A, u) = 2\partial_\mu^A f^{ia\alpha}(x_A)u_i^- + 2i(V_\mu)^\alpha{}_\beta f^{ia\beta}(x_A)u_i^-. \quad (6.23)$$

Note that the Lagrange multipliers V_μ and M_v remain, although an infinite set of auxiliary fields have been eliminated.

At this stage, we can write down the component action. Substituting the Grassmann expansions (6.14) and (6.15) into the action (6.10), using the equations of motion (6.16)–(6.19), and integrating the Grassmann variable, the action (6.10) can be put into the following form:

$$\begin{aligned} S = \int d^4 x_A du \Bigg\{ & -\bar{F}_{a\alpha}^+ \partial_A^\mu A_\mu^{-a\alpha} - i\bar{F}_{a\alpha}^+ (V^\mu)^\alpha{}_\beta A_\mu^{-a\beta} - \bar{F}_{a\alpha}^+ \tilde{m}^a_b N^{-b\alpha} \\ & + i\bar{F}_{a\alpha}^+ (\bar{M}_v)^\alpha{}_\beta N^{-a\beta} + \bar{F}_{a\alpha}^+ \tilde{m}^a_b M^{-a\alpha} + i\bar{F}_{a\alpha}^+ (M_v)^\alpha{}_\beta M^{-a\beta} \\ & + D_v^{A(-)} (i\bar{F}_{a\alpha}^+ (T_A)^\alpha{}_\beta F^{+a\beta} - \xi^{++} \text{tr}(T_A)) \Bigg\}. \end{aligned} \quad (6.24)$$

Then, using the conjugates of the equations of motion (6.17)–(6.19),

$$\bar{M}_{a\alpha}^+ = \bar{M}_{a\alpha}^i u_i^+ = \bar{F}_{b\alpha}^+ \tilde{m}^b_a + i\bar{F}_{a\beta}^+ (M_v)^\beta{}_\alpha, \quad (6.25)$$

$$\bar{N}_{a\alpha}^+ = \bar{N}_{a\alpha}^i u_i^+ = -\bar{F}_{b\alpha}^+ \tilde{m}^b_a + i\bar{F}_{a\beta}^+ (\bar{M}_v)^\beta{}_\alpha, \quad (6.26)$$

$$\bar{A}_{\mu a\alpha}^+ = \bar{A}_{\mu a\alpha}^i u_i^+ = 2(\partial_\mu^A \bar{F}_{a\alpha}^+ - i\bar{F}_{a\beta}^+ (V_\mu)^\beta{}_\alpha), \quad (6.27)$$

where $\bar{F}^+ = \bar{f}^i u_i^+$, we can rewrite the action in the simple form

$$S = \int d^4 x_A du \Bigg\{ \bar{A}_{a\alpha}^+ A_\mu^{-a\alpha} + \bar{M}_{a\alpha}^+ M^{-a\alpha} + \bar{N}_{a\alpha}^+ N^{-a\alpha} \Bigg\}$$

$$+D_v^A{}^{(-)}(i\bar{F}_{a\alpha}^+(T_A)^\alpha{}_\beta F^{+a\beta} - \xi^{(++)}\text{tr}(T_A)) \Big\}. \quad (6.28)$$

The harmonic variables can be integrated easily by using the following formulas:

$$\int du u_i^+ u_j^- = \frac{1}{2} \epsilon_{ij}, \quad (6.29)$$

$$\int du u_i^+ u_j^+ u_k^- u_l^- = \frac{1}{6} (\epsilon_{il} \epsilon_{jk} + \epsilon_{ik} \epsilon_{jl}). \quad (6.30)$$

The action after integration over u is

$$S = \int d^4 x_A (\mathcal{L}_{\text{kin}} + \mathcal{L}_{\text{constr}} + \mathcal{L}_{\text{pot}}), \quad (6.31)$$

$$\mathcal{L}_{\text{kin}} = -\frac{1}{2} A_{\mu}^{ia\alpha} \bar{A}_{ia\alpha}^{\mu}, \quad (6.32)$$

$$\mathcal{L}_{\text{constr}} = \frac{1}{3} D_{v(ij)}^A \left(i\bar{f}_{a\alpha}^{(i}(T_A)^\alpha{}_\beta f^{j)a\beta} - \xi^{(ij)} \text{tr}(T_A) \right), \quad (6.33)$$

$$\mathcal{L}_{\text{pot}} = -\frac{1}{2} (M^{ia\alpha} \bar{M}_{ia\alpha} + N^{ia\alpha} \bar{N}_{ia\alpha}) = -V. \quad (6.34)$$

After substituting (6.20)–(6.23) into (6.32)–(6.34) and integrating over the auxiliary field $D_{v(ij)}^A$, we obtain the bosonic component Lagrangian,

$$\begin{aligned} \mathcal{L}_{\text{kin}} = & -2(\partial_\mu^A f^{ia\alpha} \partial_A^\mu \bar{f}_{ia\alpha} - i\bar{f}_{ia\alpha} (V^\mu)^\alpha{}_\beta \partial_\mu^A f^{ia\beta} \\ & + i\partial_A^\mu \bar{f}_{ia\alpha} (V_\mu)^\alpha{}_\beta f^{ia\beta} + \bar{f}_{ia\alpha} (V^\mu)^\alpha{}_\beta (V_\mu)^\beta{}_\gamma f^{ia\gamma}), \end{aligned} \quad (6.35)$$

$$\begin{aligned} V = & \frac{1}{2} \bar{f}_{ia\alpha} \{\tilde{m}, \tilde{m}\}_b^a f^{ib\alpha} + \frac{1}{2} \bar{f}_{ia\alpha} \{M_v, \bar{M}_v\}_\beta^\alpha f^{ia\beta} \\ & - i\bar{f}_{ia\alpha} (\bar{M}_v)^\alpha{}_\beta \tilde{m}^a_b f^{ib\beta} + i\bar{f}_{ia\alpha} \tilde{m}^a_b (M_v)^\alpha{}_\beta f^{ia\beta}, \end{aligned} \quad (6.36)$$

with the constraint

$$i\bar{f}_{a\alpha}^{(i}(T_A)^\alpha{}_\beta f^{j)a\beta} - \xi^{(ij)} \text{tr}(T_A) = 0, \quad (6.37)$$

which makes the target space nontrivial. This constraint can also be obtained as the lowest-order component of (6.13). Let us next eliminate the Lagrange multipliers V_μ and M_v . In order to do so, we define

$$f^{ia\alpha} \bar{f}_{ia\beta} \equiv G^A(T_A)^\alpha{}_\beta, \quad i f^{ia\alpha} \overleftrightarrow{\partial}_\mu \bar{f}_{ia\beta} \equiv H_\mu^A(T_A)^\alpha{}_\beta, \quad (6.38)$$

$$\tilde{G}_{AB} \equiv \frac{1}{2} \text{tr}(\{T_A, T_B\} T_C) G^C, \quad (6.39)$$

$$\bar{f}_{ia\beta} \tilde{m}^a_b f^{ib\alpha} \equiv L^A(T_A)^\alpha{}_\beta. \quad (6.40)$$

Using these quantities, we can rewrite the Lagrangian as

$$\mathcal{L}_{\text{kin}} = -2(\partial_\mu^A f^{ia\alpha} \partial_A^\mu \bar{f}_{ia\alpha} + V_\mu^A H_A^\mu + V_\mu^A V^{B\mu} \tilde{G}_{AB}), \quad (6.41)$$

$$V = \bar{f}_{ia\alpha} |\tilde{m}|^{2a} f^{ib\alpha} + M_v^A \bar{M}_v^B \tilde{G}_{AB} + iM_v^A \bar{L}_A - i\bar{M}_v^A L_A. \quad (6.42)$$

The equations of motion for V_μ and M_v are given by

$$\frac{\partial \mathcal{L}}{\partial V_\mu^A} = -H_A^\mu - 2V^{B\mu} \tilde{G}_{AB} = 0, \quad (6.43)$$

$$\frac{\partial \mathcal{L}}{\partial M_v^A} = \bar{M}_v^B \tilde{G}_{AB} + i\bar{L}_A = 0. \quad (6.44)$$

Solving these and substituting into (6.41) and (6.42), we finally obtain

$$\mathcal{L}_{\text{kin}} = -2 \left(\partial_\mu^A f^{ia\alpha} \partial_A^\mu \bar{f}_{ia\alpha} - \frac{1}{4} H_\mu^A (\tilde{G}^{-1})_{AB} H^{B\mu} \right), \quad (6.45)$$

$$V = \bar{f}_{ia\alpha} |\tilde{m}|^{2a} f^{ib\alpha} - L^A (\tilde{G}^{-1})_{AB} \bar{L}^B. \quad (6.46)$$

The bosonic Lagrangian of the $SU(M)$ model has the same form as that in the $U(M)$ model if we regard the parameters b and c as arbitrary complex and real parameters, respectively.

6.3. Vacua in the massive $T^*G_{N,M}$ model

We now study vacua based on the potential (6.34) and the constraint. The SUSY vacuum conditions are that the auxiliary fields vanish:

$$M^{ia\alpha} = N^{ia\alpha} = 0. \quad (6.47)$$

These conditions imply vanishing values for the potential (6.34) which is clearly positive definite. We should, of course, simultaneously impose the constraint (6.37).

It is convenient to rewrite these conditions as

$$f^{ia\alpha} = \begin{pmatrix} \phi^{a\alpha} \\ -i\bar{\chi}^{a\alpha} \end{pmatrix}, \quad \bar{f}_{ia\alpha} = \epsilon^{ij} \overline{f_{ja\alpha}} = \epsilon^{ij} \bar{f}_{ja\alpha} = \begin{pmatrix} i\chi_{a\alpha} \\ -\bar{\phi}_{a\alpha} \end{pmatrix}, \quad (6.48)$$

$$\xi^{(11)} = -ib^*, \quad \xi^{(12)} = \xi^{(21)} = \frac{ic}{2}, \quad \xi^{(22)} = ib, \quad (6.49)$$

$$iM_v = \sigma. \quad (6.50)$$

In terms of these variables, Eq. (6.47) becomes

$$0 = \tilde{m}^a_b \phi^{b\alpha} + \bar{\sigma}^\alpha_\beta \phi^{a\beta}, \quad (6.51)$$

$$0 = \tilde{m}^a{}_b \bar{\chi}^{b\alpha} + \bar{\sigma}^\alpha{}_\beta \bar{\chi}^{a\beta}, \quad (6.52)$$

$$0 = \tilde{m}^a{}_b \phi^{b\alpha} + \sigma^\alpha{}_\beta \phi^{a\beta}, \quad (6.53)$$

$$0 = \tilde{m}^a{}_b \bar{\chi}^{b\alpha} + \sigma^\alpha{}_\beta \bar{\chi}^{a\beta}. \quad (6.54)$$

Also, the constraints in Eq. (6.37) become

$$0 = \chi_{\alpha a} \phi^{a\beta} = \bar{\phi}_{\alpha a} \bar{\chi}^{a\beta}, \quad (6.55)$$

$$0 = \chi_{\alpha a} \bar{\chi}^{a\beta} - \bar{\phi}_{\alpha a} \phi^{a\beta} - c \delta_\alpha{}^\beta. \quad (6.56)$$

Let us solve Eqs (6.51)–(6.56). Using $U(M)$ rotation, σ can be diagonalized as

$$\sigma = \text{diag.}(\sigma_1, \sigma, \dots, \sigma_M), \quad (6.57)$$

$$\sum_{\alpha=1}^M \sigma_\alpha = M \sigma^0. \quad (\sigma = \sigma^A T_A)$$

Equations (6.51)–(6.54) can be rewritten in terms of the diagonal masses $\tilde{m} = \text{diag.}(m_1, \dots, m_N)$ and σ as

$$0 = (\tilde{m}_a \mathbf{1}_M + \bar{\sigma}_\alpha \mathbf{1}_N) \phi^{a\alpha}, \quad (6.58)$$

$$0 = (\tilde{m}_a \mathbf{1}_M + \bar{\sigma}_\alpha \mathbf{1}_N) \bar{\chi}^{a\alpha}, \quad (6.59)$$

$$0 = (m_a \mathbf{1}_M + \sigma_\alpha \mathbf{1}_N) \phi^{a\alpha}, \quad (6.60)$$

$$0 = (m_a \mathbf{1}_M + \sigma_\alpha \mathbf{1}_N) \bar{\chi}^{a\alpha}. \quad (6.61)$$

Because $M^{ia\alpha}$ and $N^{ia\alpha}$ are not complex conjugates of each other, Eqs. (6.58) and (6.59) are not complex conjugates of Eqs. (6.60) and (6.61). However, these four conditions are consistent.

Because the SUSY conditions (6.55)–(6.59), together with the constraints (6.60) and (6.61), are the same form as (4.20)–(4.23), we can repeat the analysis given for the $\mathcal{N} = 1$ formulation in §4.3 to obtain the solutions. We thus find that there are $N C_M$ solutions of SUSY vacua.

§7. Summary and discussion

We have constructed the massive $T^*G_{N,M}$ model and its generalization, which are the Higgs branch of $\mathcal{N} = 2$ SUSY gauge theories with $U(M)$ and $SU(M)$ gauge groups, respectively. The vacuum structure for the massive $T^*G_{N,M}$ model has been

clarified, and it was found to be far richer than the massive $T^*\mathbf{CP}^{N-1}$ model. It has discrete $N!/ [M!(N-M)!]$ vacua, which are the origins of the standard coordinate patches for $G_{N,M}$ and are represented by mutually orthogonal M -dimensional complex planes in $\mathbf{C}^N$ in a coordinate-independent way. On the other hand, the massive HK sigma model with the $SU(M)$ quotient has no discrete vacua.

Since we have discrete vacua for the massive $T^*G_{N,M}$ model, we believe that this model has domain wall configurations that are richer than the massive $T^*\mathbf{CP}^{N-1}$ model. First, even for a single wall, we can expect several types of solutions. For example, the cotangent bundle over the Klein quadric, $T^*G_{4,2}$, has the six vacua given in Eq. (3.14). This model is expected to admit two kinds of walls. One is a wall connecting two vacua without any common axis, e.g., the first and the sixth vacua. The other is a wall connecting two vacua with a common axis, e.g., the first and the second vacua. For general massive $T^*G_{N,M}$ models, we conjecture a greater variety of wall solutions. Second, if we have a parallel wall configuration, as in the massive $T^*\mathbf{CP}^{N-1}$ model, such a configuration has several zero modes corresponding to positions of the walls.^{13), 27)} The number of zero modes was calculated to be N for the massive $T^*\mathbf{CP}^{N-1}$ model in Ref. 29) using the index theorem. It is interesting to apply the approach used there to the massive $T^*G_{N,M}$ model.

The second homotopy group for $T^*G_{N,M}$ is nontrivial: $\pi_2(T^*G_{N,M}) \simeq \pi_2(G_{N,M}) \simeq \mathbf{Z}$.⁴⁹⁾ Therefore, this model is believed to admit multi-string (vortex or lump) solutions, a configuration of a string ending on a wall, as was found in Ref. 31) for $T^*\mathbf{CP}^1$, and other interesting phenomena.

Reducing the $T^*\mathbf{CP}^{N-1}$ model to three-dimensional space-time, an interesting mirror symmetry has been found.⁵⁰⁾ Determining the non-Abelian generalization of this mirror symmetry using our massive $T^*G_{N,M}$ model would be an interesting task.

Coupling the model to supergravity is possible, and in that case, the target manifold is the quaternionic generalization of $T^*G_{N,M}$. Moreover, our model can be promoted to five-dimensional supergravity in the manner considered in Refs. 12) and 13). This is interesting for the brane world scenario.

Acknowledgements

We thank Masashi Naganuma for collaboration in the early stages of this work. This work is supported in part by a Grant-in-Aid for Scientific Research from the Japan Ministry of Education, Culture, Sports, Science and Technology 13640269 (N.S.). The work of M. N. is supported by the U. S. Department of Energy under grant DE-FG02-91ER40681 (Task B).

References

- 1) N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *Phys. Lett.* **B429** (1998) 263, [hep-ph/9803315]; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *Phys. Lett.* **B436** (1998) 257, [hep-ph/9804398].
- 2) L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83** (1999) 3370, [hep-ph/9905221]; *Phys. Rev. Lett.* **83** (1999) 4690, [hep-th/9906064].
- 3) S. Dimopoulos and H. Georgi, *Nucl. Phys.* **B193** (1981) 150; N. Sakai, *Z. f. Phys.* **C11** (1981) 153; E. Witten, *Nucl. Phys.* **B188** (1981) 513; S. Dimopoulos, S. Raby and F. Wilczek, *Phys. Rev.* **D24** (1981) 1681.
- 4) E. Bogomol'nyi, *Sov. J. Nucl. Phys.* **B24** (1976) 449; M.K. Prasad and C.H. Sommerfield, *Phys. Rev. Lett.* **35** (1975) 760.
- 5) E. Witten and D. Olive, *Phys. Lett.* **B78** (1978) 97.
- 6) N. Seiberg and E. Witten, *Nucl. Phys.* **B426** (1994) 19, [hep-th/9407087]; *Nucl. Phys.* **B431** (1994) 484, [hep-th/9408099].
- 7) P.C. Argyres, M.R. Plesser and N. Seiberg, *Nucl. Phys.* **B471** (1996) 159, [hep-th/9603042].
- 8) I. Antoniadis and B. Pioline, *Int. J. Mod. Phys.* **A12** (1997) 4907, [hep-th/9607058].
- 9) M. Shifman and A. Yung, *Phys. Rev.* **D67** (2003) 125007, [hep-th/0212293].
- 10) K. Evlampiev and A. Yung, *Nucl. Phys.* **B662** (2003) 120, [hep-th/0303047].
- 11) M. Arai, M. Naganuma, M. Nitta and N. Sakai, *Nucl. Phys.* **B652** (2003) 35, [hep-th/0211103]; “BPS Wall in N=2 SUSY Nonlinear Sigma Model with Eguchi-Hanson Manifold” in Garden of Quanta - In honor of Hiroshi Ezawa, Eds. by J. Arafune et al. (World Scientific Publishing Co. Pte. Ltd. Singapore, 2003) pp 299-325, [hep-th/0302028].
- 12) M. Arai, S. Fujita, M. Naganuma and N. Sakai, *Phys. Lett.* **B556** (2003) 192, [hep-th/0212175].
- 13) M. Eto, S. Fujita, M. Naganuma and N. Sakai, *Phys. Rev.* **D69** (2004) 025007, [hep-th/0306198].
- 14) B. Zumino, *Phys. Lett.* **B87** (1979) 203.
- 15) L. Alvarez-Gaumé and D.Z. Freedman, *Comm. Math. Phys.* **80** (1981) 443.
- 16) L. Alvarez-Gaumé and D.Z. Freedman, *Comm. Math. Phys.* **91** (1983) 87.
- 17) J. Scherk and J.H. Schwarz, *Phys. Lett.* **B82** (1979) 60.
- 18) G. Sierra and P.K. Townsend, *Nucl. Phys.* **B233** (1984) 289.

- 19) J.Gates, Jr., *Nucl. Phys.* **B238** (1984) 349.
- 20) A. Losev and M. Shifman *Phys. Rev.* **D68** (2003) 045006, [hep-th/0304003].
- 21) K. Kakimoto and N. Sakai, *Phys. Rev.* **D68** (2003) 065005, [hep-th/0306077].
- 22) U. Lindström and M. Roček, *Nucl. Phys.* **B222** (1983) 285.
- 23) H. Pedersen and Y.S. Poon, *Comm. Math. Phys.* **117** (1988) 569.
- 24) G.W. Gibbons, R. Goto and P. Rychenkova, *Comm. Math. Phys.* **185** (1997) 581, [hep-th/9608085].
- 25) R. Bielawski, *Math. Ann.* **314** (1999) 505, [math. DG/9808134].
- 26) J.P. Gauntlett, D. Tong and P.K. Townsend, *Phys. Rev.* **D63** (2001) 085001, [hep-th/0007124].
- 27) J.P. Gauntlett, D. Tong and P.K. Townsend, *Phys. Rev.* **D64** (2001) 025010, [hep-th/0012178].
- 28) D. Tong, *Phys. Rev.* **D66** (2002) 025013, [hep-th/0202012].
- 29) K.S.M. Lee, *Phys. Rev.* **D67** (2003) 045009, [hep-th/0211058].
- 30) E. Abraham and P.K. Townsend, *Phys. Lett. B* **291** (1992) 85.
- 31) J.P. Gauntlett, R. Portugues, D. Tong and P.K. Townsend, *Phys. Rev.* **D63** (2001) 085002, [hep-th/0008221].
- 32) M. Naganuma, M. Nitta and N. Sakai, *Grav. Cosmol.* **8** (2002) 129, [hep-th/0108133].
- 33) R. Portugues and P.K. Townsend, *JHEP* **0204** (2002) 039, [hep-th/0203181].
- 34) A. Hanany and K. Hori, *Nucl. Phys.* **B513** (1998) 119, [hep-th/9707192].
- 35) N. J. Hitchin, A. Karlhede, U. Lindström and M. Roček, *Comm. Math. Phys.* **108** (1987) 535.
- 36) A. Galperin, E. Ivanov, S. Kalitzin, V. Ogievetsky and E. Sokatchev, *Class. Quantum Grav.* **1** (1984) 469; A. Galperin, E. Ivanov, V. Ogievetsky and E. Sokatchev, *Class. Quantum Grav.* **2** (1985) 601; 617.
- 37) A. Galperin, E. Ivanov, V. Ogievetsky and E. Sokatchev, “Harmonic Superspace”, Cambridge University Press, Cambridge (2001).
- 38) K. Higashijima and M. Nitta, *Prog. Theor. Phys.* **103** (2000) 635, [hep-th/9911139].
- 39) T.L. Curtright and D.Z. Freedman, *Phys. Lett.* **B90** (1980) 71.
- 40) L. Alvarez-Gaumé and D.Z. Freedman, *Phys. Lett.* **B94** (1980) 171.
- 41) M. Roček and P.K. Townsend, *Phys. Lett.* **B96** (1980) 72.
- 42) E. Calabi, *Ann. Scient. Ec. Norm. Sup.* **12** (1979) 269.
- 43) T. Eguchi and A. J. Hanson, *Phys. Lett.* **B74** (1978) 249; *Ann. Phys.* **120** (1979) 82.
- 44) K. Higashijima, T. Kimura and M. Nitta, *Nucl. Phys.* **B623** (2002) 133, [hep-th/0108084]; *Annals Phys.* **296** (2002) 347, [hep-th/0110216].
- 45) M. Naganuma, M. Nitta and N. Sakai, *Phys. Rev.* **D65** (2001) 045016, [hep-th/0108179]; “BPS Walls and Junctions in $\mathcal{N} = 1$ SUSY Nonlinear Sigma Models”, The Proceedings of the third international Sakharov conference on Physics, Vol.2, (2003) p535-549, World Scientific Pub., [hep-th/0210205].
- 46) A. Galperin, E. Ivanov, V. Ogievetsky and P.K. Townsend, *Class. Quantum Grav.* **3** (1986) 625.
- 47) A. Galperin, E. Ivanov, V. Ogievetsky and E. Sokatchev, *Comm. Math. Phys.* **103** (1986)

515.

48) S. Ketov and C. Unkmeir, *Phys. Lett.* **B422** (1998) 179, [hep-th/9710185].

49) A.M. Perelomov, *Phys. Rept.* **146** (1987) 135.

50) D. Tong, *JHEP* **0304** (2003) 031, [hep-th/0303151].